

CLASSIFICATION OF GORENSTEIN TORIC DEL PEZZO VARIETIES IN ARBITRARY DIMENSION

VICTOR BATYREV AND DOROTHEE JUNY

Dedicated to the memory of Professor V. A. Iskovskih

ABSTRACT. A n -dimensional Gorenstein toric Fano variety X is called Del Pezzo variety if the anticanonical class $-K_X$ is a $(n-1)$ -multiple of a Cartier divisor. Our purpose is to give a complete biregular classification of Gorenstein toric Del Pezzo varieties in arbitrary dimension $n \geq 2$. We show that up to isomorphism there exist exactly 37 Gorenstein toric Del Pezzo varieties of dimension n which are not cones over $(n-1)$ -dimensional Gorenstein toric Del Pezzo varieties. Our results are closely related to the classification of all Minkowski sum decompositions of reflexive polygons due to Emiris and Tsigaridas and to the classification up to deformation of n -dimensional almost Del Pezzo manifolds obtained by Jahnke and Peternell.

INTRODUCTION

A projective algebraic variety X is called *Gorenstein Fano variety* if the anticanonical divisor $-K_X$ is an ample Cartier divisor. A n -dimensional Gorenstein Fano variety is called *Gorenstein Del Pezzo variety* if

$$-K_X = (n-1)L$$

for some ample Cartier divisor L on X . Smooth Del Pezzo n -folds have been classified up to deformation by Iskovskih and Fujita [Isk77, Isk78, Isk80, Fuj80, Fuj81, Fuj84]. A classification of Gorenstein Del Pezzo surfaces is known due to Demazure [Dem80].

All Gorenstein Del Pezzo varieties of dimension $n \geq 4$ have been classified up to deformation by Fujita [Fuj90a, Fuj90b]. A smooth n -fold X is called an *almost Del Pezzo manifold* if there exists a K_X -trivial birational morphism $\varphi : X \rightarrow X'$ where X' is a Gorenstein Del Pezzo variety. It is well-known that if $n = 2$ then the minimal resolution $\hat{X}$ of a Gorenstein Del Pezzo surface X is always an almost Del Pezzo surface [DuV34]. Such a surface $\hat{X}$ can be obtained from $\mathbb{P}^2$ or $\mathbb{P}^1 \times \mathbb{P}^1$ by blowing up points which are not necessary in general position. Recently all almost Del Pezzo manifolds of arbitrary dimension n were classified up to deformation by Jahnke and Peternell [JP08].

In this paper we consider n -dimensional Gorenstein Fano varieties X which are *toric varieties*, i.e., X contains a n -dimensional algebraic torus T as an open dense subset such that the group multiplication $T \times T \rightarrow T$ extends to a regular action $T \times X \rightarrow X$ [Oda88, Ful93]. Let $M \cong \mathbb{Z}^n$ be the lattice of characters of T and $M_{\mathbb{R}} := M \otimes \mathbb{R}$ the corresponding real vector space. A natural T -action on the space

$H^0(X, \mathcal{O}(-K_X))$ splits into a finite direct sum of 1-dimensional weight subspaces parametrized by a finite subset of characters $\{\chi_i\} \subset M$. In this situation, the convex hull Δ of the set $\{\chi_i\}$ is a special n -dimensional polytope in $M_{\mathbb{R}}$ which is called *reflexive polytope* [Bat94]. The Gorenstein toric Fano variety X is uniquely determined up to an isomorphism of the n -dimensional lattice M by the corresponding reflexive polytope Δ . It is known that for any given dimension n there exist only finitely many n -dimensional reflexive polytopes up to a lattice isomorphism. This implies that there exist only finitely many Gorenstein toric Fano varieties of dimension n up to biregular isomorphism (see [Bat82]). If $N := \text{Hom}(M, \mathbb{Z})$ is the dual lattice and $N_{\mathbb{R}} = \text{Hom}(M_{\mathbb{R}}, \mathbb{R})$ the dual real vector space, then for every reflexive polytope $\Delta \subset M_{\mathbb{R}}$ there exists a dual reflexive polytope $\Delta^* \subset N_{\mathbb{R}}$. Using an isomorphism $N \cong M$, we obtain that up to isomorphism all reflexive polytopes satisfy a nice combinatorial duality which plays important role in Mirror Symmetry [Bat94] (see also [CK99]).

If X is a n -dimensional toric variety such that $-K_X = rL$ for an ample divisor L then, up to a translation by an element of M , the reflexive polytope Δ is isomorphic to a r -multiple of another lattice polytope P , i.e., $\Delta \cong rP$ where all vertices of P are contained in M . Such polytopes P are called *Gorenstein polytopes of index r* [BR07, BN08]. For example, the standard n -dimensional basic lattice simplex is a Gorenstein polytope of index $n + 1$ and the standard n -dimensional lattice unit cube is a Gorenstein polytope of index 2. The number $d := n + 1 - r$ is called the *degree* of a n -dimensional Gorenstein polytope P (see also [Bat06]). The name “degree” of a n -dimensional lattice polytope P is motivated by the following fact: if

$$E_P(t) := \sum_{k=0}^{\infty} |kP \cap M| t^k$$

is the Ehrhart series of P , then $h_P^*(t) := (1 - t)^{n+1} E_P(t)$ is a polynomial of degree d .

The purpose of our paper is to give a complete classification up to isomorphism of all Gorenstein toric Del Pezzo varieties in arbitrary dimension $n \geq 2$. This is equivalent to a classification of all n -dimensional Gorenstein polytopes of index $r = n - 1$, or, equivalently, of degree 2. In the case $n = 2$, Gorenstein polygons of index 1 are exactly reflexive polygons and their classification is well-known (it was obtained independently in [Bat85, Rab89, Koe91]). There exist exactly 16 reflexive polygons (see Figure 1).

It is important to note that if X is a n -dimensional Gorenstein Del Pezzo variety then the cone X' over X is a $(n + 1)$ -dimensional Gorenstein Del Pezzo variety. If in addition X is toric, then X' is also toric and the $(n + 1)$ -dimensional Gorenstein polytope P' corresponding to X' is a lattice pyramid $\Pi(P)$ over the n -dimensional Gorenstein polytope P corresponding to X . This shows that it is enough to classify only those Gorenstein polytopes of index $r = n - 1$ which are not pyramids over low dimensional ones.




FIGURE 1. All 16 reflexive polygons, up to isomorphism

Using a computer program, 3- and 4-dimensional reflexive polytopes have been classified by Kreuzer and Skarke [KS98, KS00]. Their computer calculation resulted in 4319 polytopes in dimension 3 and 473 800 776 polytopes in dimension 4. Using the lists of 3- and 4-dimensional polytopes, Kreuzer has found 31 3-dimensional Gorenstein polytopes of index 2 and 36 4-dimensional Gorenstein polytopes of index 3. In this paper, we give an independent proof of this fact without using computer classifications.

Our complete classification of n -dimensional Gorenstein polytopes of degree 2 (i.e., of index $r = n - 1$) gives the following more precise result:

Dimension n	Number of polytopes	Number of non-pyramids
2	16	16
3	31	15
4	36	5
5	37	1
≥ 6	37	-

We remark that the classification of n -dimensional Gorenstein polytopes of index $r = n - 2$ is a much harder problem. This problem in case $n = 3$ is equivalent to the classification of all 3-dimensional reflexive polytopes [KS98]. Using the computer classification of 4-dimensional reflexive polytopes [KS00], Kreuzer has found exactly

5363 reflexive polytopes Δ of dimension 4 which are isomorphic to $2P$ for some lattice polytope P . This shows that there exist exactly 5363 Gorenstein polytopes P of dimension 4 and index 2. In particular, there exist exactly $1044 = 5363 - 4319$ Gorenstein 4-dimensional polytopes of index 2 which are not pyramids over 3-dimensional reflexive polytopes. However, all these facts seem to be very difficult to check without computer.

It follows from a recent result of Haase, Nill and Payne [HNP08, Cor. 3.2] together with the main result in [Bat06] that every n -dimensional Gorenstein polytope P of degree d is a pyramid if

$$n \geq 4d \binom{2d + V - 1}{2d},$$

where

$$V = (2d - 1)^{2d-1} \cdot ((2d - 1)!)^{2d} \cdot 14^{(2d-1)^2 \cdot 2^{2d}}.$$

In [Nil08, Prop. 1.5] Nill has shown that the same statement holds true already for

$$n \geq (V - 1)(2d + 1).$$

However, these inequalities are still too far from being sharp. It would be nice to know exactly the maximal dimension n (depending on d) such that every n -dimensional Gorenstein polytope of degree d is a pyramid. In this connection we suggest the following:

Conjecture 0.1. *Every 3d-dimensional Gorenstein polytope P of degree d is a pyramid. Moreover, there exists up to an isomorphism a unique $(3d - 1)$ -dimensional Gorenstein polytope Θ_d of degree d which is not a pyramid.*

Conjecture 0.1 is easy to check for $d = 1$. Our classification verifies Conjecture 0.1 in the case $d = 2$, but it remains open for $d \geq 3$. A precise description of the special Gorenstein polytope Θ_d for arbitrary degree d is given in 1.22.

Let us summarize the ideas used in our classification of n -dimensional Gorenstein polytopes of degree 2.

First of all we use a natural partial order on the set of n -dimensional Gorenstein polytopes of degree 2: for two lattice polytopes P_1 and P_2 , we write $P_1 \preceq P_2$ if P_1 is isomorphic to a lattice subpolytope of P_2 . We say that a n -dimensional Gorenstein polytope P of degree 2 is *minimal* if there does not exist a n -dimensional Gorenstein subpolytope $P' \preceq P$ of degree 2 such that $P' \neq P$. For n -dimensional Gorenstein polytopes of fixed degree d there exists a combinatorial duality which generalize the polar duality for n -dimensional reflexive polytopes [BN08]. Using this duality, we obtain that if P is a minimal n -dimensional Gorenstein polytope of degree 2, then the dual Gorenstein polytope P^* is a *maximal* one, i.e., P^* is not contained in a strictly larger n -dimensional Gorenstein polytope of degree 2. Since every n -dimensional Gorenstein polytope is contained in a maximal one, in the classification of all n -dimensional Gorenstein polytopes of degree 2 the first step is to find all minimal

ones among them. After that, using the duality $P \leftrightarrow P^*$, we immediately obtain all maximal n -dimensional Gorenstein polytopes of degree 2. Finally, all remaining Gorenstein polytopes can be found as lattice subpolytopes of the maximal ones.

Let P be a n -dimensional lattice polytope. The integer

$$\text{Vol}_{\mathbb{N}}(P) := n!(\text{volume of } P)$$

we call the *lattice normalized volume of P* . In the classification of minimal Gorenstein polytopes of degree 2 the following statement is very important: if P is minimal, then $\text{Vol}_{\mathbb{N}}(P) \leq 4$. The proof of this technical statement consists of several steps and uses the complete classification of n -dimensional lattice polytopes whose h^* -polynomial is of degree ≤ 1 [BN07]. As we already remarked, the classification of all n -dimensional Gorenstein polytopes P of degree 2 is equivalent to the classification of all n -dimensional lattice polytopes P having the Ehrhart series

$$E_P(t) = \frac{1 + a_P t + t^2}{(1 - t)^{n+1}}, \quad a_P \in \mathbb{Z}_{\geq 0},$$

where $a_P = \text{Vol}_{\mathbb{N}}(P) - 2$. If X_P is the corresponding n -dimensional Gorenstein toric Del Pezzo variety and $-K_{X_P} = (n - 1)L$ for some Cartier divisor L , then the integer $a_P + 2 = \text{Vol}_{\mathbb{N}}(P)$ equals the intersection number L^n and the Ehrhart series $E_P(t)$ is exactly the Hilbert-Poincaré series of the homogeneous coordinate ring of X_P :

$$S_P := \bigoplus_{k=0}^{\infty} H^0(X_P, \mathcal{O}(kL)).$$

The graded Gorenstein ring S_P can be seen as a semigroup algebra of the set of lattice points in the $(n + 1)$ -dimensional cone C_P over the Gorenstein polytope P . If P is a Gorenstein polytope such that $\text{Vol}_{\mathbb{N}}(P) \leq 4$ (i.e. $a_P \leq 2$), then one can apply algebraic ideas from [Bat06] and classify all possible Gorenstein rings S_P (and the corresponding Gorenstein polytopes P) by an enumeration of all binomial relations among the minimal generators of S_P .

Let us say some words about the way of presentation of the results of our classification. It is not difficult to draw pictures of all 3-dimensional Gorenstein polytopes of degree 2. However, the same way of presentation can not be applied in dimension $n \geq 4$. It was shown in [BN08, Th. 2.6] that if a d -dimensional reflexive polytope Δ is a Minkowski sum of r lattice polytopes $\Delta_1, \dots, \Delta_r$:

$$\Delta = \Delta_1 + \dots + \Delta_r,$$

then the $(d + r - 1)$ -dimensional Cayley polytope $P := \Delta_1 * \dots * \Delta_r$ is a Gorenstein polytope of degree d (or, of index r). In particular, a Minkowski sum decomposition $\Delta = \Delta_1 + \dots + \Delta_r$ of a 2-dimensional reflexive polytope Δ defines a $(r + 1)$ -dimensional Gorenstein polytope $P = \Delta_1 * \dots * \Delta_r$ of degree 2. The classification of all Minkowski sum decompositions of reflexive polygons was recently obtained by Emiris and Tsirigaris [ET06]. Our classification shows that all Gorenstein polytopes of degree 2 and of dimension $n \geq 4$ are Cayley polytopes corresponding to different Minkowski

sum decompositions of 2-dimensional reflexive polytopes. The same statement holds true for almost all (except one) 3-dimensional Gorenstein polytopes of degree 2.

The paper is organized as follows:

In Section 1 we review basic definitions and properties of Gorenstein polytopes. The combinatorial duality of Gorenstein polytopes $P \leftrightarrow P^*$ and their constructions as Cayley polytopes corresponding to Minkowski summands of reflexive polygons will be of our primary interest.

In Section 2 we investigate the graded Gorenstein K -algebra S_P associated with a Gorenstein polytope P of degree d . Let $A_P := S_P/\langle y \rangle$ be the Artinian Gorenstein K -algebra obtained from S_P as quotient modulo an ideal $\langle y \rangle$ generated by a regular sequence of homogeneous elements $y_1, \dots, y_n$ of degree 1. If the degree of P is 2 then $1 + a_P t + t^2$ is the Hilbert-Poincaré polynomial of A_P and the graded Gorenstein K -algebras A_P can be classified up to isomorphism (see 2.9). This classification allows to determine all possible binomial relations in the K -algebra S_P . If $a_P \leq 2$, then we find a complete list of Gorenstein polytopes P which are not pyramids (see 2.10).

In Section 3 we describe general combinatorial properties of Gorenstein polytopes of degree 2. In particular, we prove some useful equalities for 3-dimensional Gorenstein polytopes of degree 2. It is well known that if Δ and Δ^* are dual to each other reflexive polygons then

$$\mathrm{Vol}_{\mathbb{N}}(\Delta) + \mathrm{Vol}_{\mathbb{N}}(\Delta^*) = 12.$$

This remarkable property of reflexive polygons was recently investigated by many mathematicians [PR00, HS02, RST05]. We show that there exists a similar relation between the number 12 and dual to each other 1-dimensional faces $E \subset P$, $E^* \subset P^*$ of the 3-dimensional Gorenstein polytopes P , P^* of degree 2

$$\sum_{E \subset P, \dim E=1} \mathrm{Vol}_{\mathbb{N}}(E) \cdot \mathrm{Vol}_{\mathbb{N}}(E^*) = 12.$$

(see 3.4).

In Section 4 we prove that if a Gorenstein polytope P of degree 2 is minimal then $\mathrm{Vol}_{\mathbb{N}}(P) \leq 4$. As it was explained above, together with the classification in Section 2 this result easily implies a complete classification of Gorenstein polytopes of degree P . Here we present the whole list of d -dimensional Gorenstein polytopes of degree 2 which are not pyramids (including some of their combinatorial properties).

In Section 5 we compare our combinatorial method with the birational method of Jahnke and Peternell used in their classification of almost Del Pezzo manifolds [JP08].

Acknowledgements. We are very grateful to Jürgen Hausen, Priska Jahnke and Benjamin Nill for their interest in our work and helpful comments.

1. GENERAL PROPERTIES OF GORENSTEIN POLYTOPES

By a *lattice* M of dimension n we mean a free abelian group of rank n , $M \cong \mathbb{Z}^n$. We also consider the *dual lattice* $N = M^* = \text{Hom}(M, \mathbb{Z}) \cong \mathbb{Z}^n$ together with the canonical pairing

$$\langle \cdot, \cdot \rangle : M \times N \rightarrow \mathbb{Z}.$$

Further, we define $M_{\mathbb{Q}} := M \otimes_{\mathbb{Z}} \mathbb{Q}$, $N_{\mathbb{Q}} := N \otimes_{\mathbb{Z}} \mathbb{Q}$ together with the $\mathbb{Q}$ -bilinear pairing $\langle \cdot, \cdot \rangle : M_{\mathbb{Q}} \times N_{\mathbb{Q}} \rightarrow \mathbb{Q}$. The real vector spaces $M_{\mathbb{R}}$ and $N_{\mathbb{R}}$ are defined analogously. A *polytope* P in $M_{\mathbb{R}}$ is the convex hull of a finite subset in $M_{\mathbb{R}}$. We say that P is a *lattice polytope* if all its vertices lie in M . Further, a *face* of P is a subset $F \subseteq P \subset M_{\mathbb{R}}$ which minimizes on P some linear function $f \in N_{\mathbb{R}}$, and a *facet* of P is a codimension-1 face of P . By ∂P we denote the boundary of the polytope P and by $\text{int}(P)$ the interior $P \setminus \partial P$ of P .

Let P be a n -dimensional lattice polytope in $M_{\mathbb{R}}$. For any nonnegative integer k the number $i(P, k) := |kP \cap M|$ is a polynomial of degree n in k [Ehr67]. This polynomial is called the *Ehrhart polynomial* of P . Its values in negative integers are defined by the formula

$$i(P, -k) = (-1)^n |\text{int}(kP) \cap M|.$$

Moreover, one has

$$E_P(t) := \sum_{k \geq 0} i(P, k) t^k = \frac{h_0^* + h_1^* t + \cdots + h_n^* t^n}{(1-t)^{n+1}}$$

for some nonnegative integers h_i^* [Sta80]. The Ehrhart polynomial $i(P, k)$ can be recovered from the integers h_i^* by the formula:

$$i(P, k) = \sum_{j=0}^n h_j^* \binom{n+k-j}{n}.$$

Definition 1.1. The polynomial

$$h_P^*(t) := (1-t)^{n+1} E_P(t) = \sum_{k \geq 0} |kP \cap M| t^n$$

is called the *h^* -polynomial* of a n -dimensional lattice polytope P . Its degree is called the *degree of P* : $\deg P := \max\{i : h_i^* \neq 0\}$, $0 \leq \deg P \leq \dim P$. The vector $h^*(P) = (h_0^*, h_1^*, \dots, h_{\deg P}^*)$ is called the *h^* -vector* of P .

Remark 1.2. It was proved by Stanley in [Sta80] that the numbers $h_i^* = h_i^*(P)$ are monotone functions of P : if P and P' are two lattice polytopes and $P' \subseteq P$, then $h_i^*(P') \leq h_i^*(P)$ for all i .

Definition 1.3. [BN07, Def. 2.1] We call a n -dimensional lattice polytope P ($n \geq 1$) a *Lawrence prism with heights* $\theta_1, \dots, \theta_n$ if there exists a lattice basis $e_1, \dots, e_n$ of M and non-negative integers $\theta_1, \dots, \theta_n$ such that

$$P = \text{conv}\{0, \theta_1 e_n, e_1, e_1 + \theta_2 e_n, \dots, e_{n-1}, e_{n-1} + \theta_n e_n\}.$$

The lattice vector e_n is called the *direction of P* .

Definition 1.4. [BN07, Def. 2.2] We call a n -dimensional lattice polytope P ($n \geq 2$) *exceptional* if there exists a lattice basis $e_1, \dots, e_n$ of M such that

$$P = \text{conv}\{0, 2e_1, 2e_2, e_3, \dots, e_n\}.$$

The following statement was proved in [BN07]:

Theorem 1.5. *Every n -dimensional lattice polytope P of degree ≤ 1 is either a basic simplex, or an exceptional simplex, or a Lawrence prism such that $\theta_1 + \dots + \theta_n \geq 2$. The corresponding h^* -polynomial is equal respectively to 1, $1+3t$, $1+(\theta_1 + \dots + \theta_n - 1)t$.*

Definition 1.6. For any r convex polytopes $\Delta_1, \dots, \Delta_r$ in $M_{\mathbb{R}}$, their *Minkowski sum* is defined as

$$\Delta := \Delta_1 + \dots + \Delta_r = \{x_1 + \dots + x_r \mid x_i \in \Delta_i\}.$$

The polytopes Δ_i ($i = 1, \dots, r$) are called *Minkowski summands* of Δ . Here we do not require all polytopes $\Delta_1, \dots, \Delta_r$ to have the maximal dimension $n = \dim M_{\mathbb{R}}$. Define the lattice $\widetilde{M} := M \oplus \mathbb{Z}^r = M \oplus \mathbb{Z}e_1 \oplus \dots \oplus \mathbb{Z}e_r$, where $\{e_1, \dots, e_r\}$ form a lattice basis of $\mathbb{Z}^r$. The *Cayley polytope* $\Delta_1 * \dots * \Delta_r$ associated with $\Delta_1, \dots, \Delta_r$ is the convex hull of $(\Delta_1, e_1), \dots, (\Delta_r, e_r)$ in $\widetilde{M}_{\mathbb{R}}$.

Definition 1.7. Let $\Delta \subset M_{\mathbb{R}}$ be a lattice polytope of dimension n . We set $\Delta_0 := \Delta$ and choose r 0-dimensional lattice polytopes $\Delta_1, \dots, \Delta_r$ (points). Then the polytope $\Delta_0 * \Delta_1 * \dots * \Delta_r$ is called *r -fold pyramid over Δ* and we denote it by $\Pi^r(\Delta)$. If $r = 1$ we write simply $\Pi(\Delta)$ and call $\Pi(\Delta)$ *pyramid over Δ* .

Remark 1.8. A Lawrence prism with heights $\theta_1, \dots, \theta_n$ can be considered as Cayley polytope of n 1-dimensional polytopes $[0, \theta_1], \dots, [0, \theta_n] \subset \mathbb{R}$. A n -dimensional exceptional simplex is the $(n-2)$ -fold pyramid over the lattice triangle $\text{conv}\{0, 2e_1, 2e_2\}$.

Definition 1.9. Consider a d -dimensional convex polytope Δ in $M_{\mathbb{R}}$ that contains the zero point $0 \in M_{\mathbb{R}}$ in its interior. The polytope

$$\Delta^* = \{y \in N_{\mathbb{R}} \mid \langle x, y \rangle \geq -1 \quad \forall x \in \Delta\}$$

is called *dual polytope* of Δ .

A d -dimensional lattice polytope Δ in $M_{\mathbb{R}}$ is called *reflexive* if $0 \in \Delta$ and the dual polytope Δ^* is again a lattice polytope. If Δ contains a single interior lattice point m such that $\Delta - m$ is reflexive, we say that Δ is *reflexive with respect to m* .

Remark 1.10. If Δ is reflexive polytope, then the dual polytope Δ^* is again reflexive and $(\Delta^*)^* = \Delta$. The duality $\Delta \leftrightarrow \Delta^*$ is very important for Mirror Symmetry [Bat94, CK99].

Proposition 1.11. [Hib92] *Let $\Delta \subset M_{\mathbb{R}}$ be a lattice polytope. Then Δ is reflexive if and only if for every nonnegative integer k the number of lattice points in $k\Delta$ equals the number of lattice points in the interior $\text{int}((k+1)\Delta)$ of Δ .*

Corollary 1.12. *A n -dimensional lattice polytope Δ is reflexive if and only if*

$$h_i^*(\Delta) = h_{n-i}^*(\Delta), \quad 0 \leq i \leq \frac{n}{2}.$$

Definition 1.13. Let r be a positive integer. A n -dimensional lattice polytope P is called *Gorenstein of index r* if rP is a reflexive polytope with respect to some interior lattice point $m \in \text{int}(rP)$.

Analogous to 1.11, one obtains:

Proposition 1.14. *Let $P \subset M_{\mathbb{R}}$ be a n -dimensional lattice polytope. Then P is a Gorenstein polytope of index r if and only if for every nonnegative integer k the number of lattice points in kP equals the number of lattice points in the interior $\text{int}((k+r)P)$ of Δ . In this case, one has $\deg P = n + 1 - r$.*

Corollary 1.15. *Let P be a n -dimensional lattice polytope with h^* -polynomial $h_P^*(t) = \sum_i h_i^* t^i$ and let $d = \deg P$. Then P is a Gorenstein polytope of degree $d = n + 1 - r$ if and only if*

$$h_i^*(P) = h_{d-i}^*(P), \quad 0 \leq i \leq \frac{d}{2}.$$

Definition 1.16. [BB97] Consider two $(n+1)$ -dimensional lattices $\overline{M} := M \oplus \mathbb{Z}$ and $\overline{N} := N \oplus \mathbb{Z}$ which are dual to each other with respect to the natural extension of the pairing $\langle *, * \rangle : M \times N \rightarrow \mathbb{Z}$ to $\overline{M} \times \overline{N} \rightarrow \mathbb{Z}$. For the latter we use the same notation $\langle *, * \rangle$. Let $\overline{M}_{\mathbb{R}}$ and $\overline{N}_{\mathbb{R}}$ be the corresponding real vector spaces.

A strongly convex $(n+1)$ -dimensional cone $C \in \overline{M}_{\mathbb{R}}$ with the vertex $0 = -C \cap C$ is called *Gorenstein cone* if there exists an element $n_C \in \overline{N}_{\mathbb{R}}$ such that $\langle e_i, n_C \rangle = 1$ for all primitive lattice generators e_i of C . Note that n_C is uniquely determined.

If the cone C is Gorenstein, then the set

$$P_C = \{x \in C \mid \langle x, n_C \rangle = 1\}$$

is a convex n -dimensional lattice polyhedron which is called *support* of C .

On the other hand, if P is a n -dimensional lattice polytope in $M_{\mathbb{R}}$, then

$$C_P := \{(\lambda x, \lambda) : \lambda \in \mathbb{R}_{\geq 0}, x \in P\} \subset \overline{M}_{\mathbb{R}}$$

is a Gorenstein cone with the support polytope P (here we have $n_{C_P} = (0, \dots, 0, 1)$).

Definition 1.17. [BB97] Define the *dual cone* as

$$C^{\vee} := \{y \in \overline{N}_{\mathbb{R}} : \langle x, y \rangle \geq 0 \quad \forall x \in C\}.$$

A $(n+1)$ -dimensional Gorenstein cone C is called *reflexive* if its dual cone $C^{\vee}$ is Gorenstein as well. In this case we denote by $m_{C^{\vee}} \in M$ the lattice point such that the hyperplane $\langle m_{C^{\vee}}, y \rangle$ contains all lattice points generating the dual cone $C^{\vee}$. For a reflexive Gorenstein cone, the uniquely defined positive integer

$$r_C = \langle m_{C^{\vee}}, n_C \rangle$$

is called *index* of C .

Remark 1.18. It is very important for our purpose that if C is a reflexive Gorenstein cone of index r then $C^\vee$ is also a reflexive Gorenstein cone of index r . Moreover, a Gorenstein cone is reflexive of index r if and only if its support is a Gorenstein polytope of index r (see [BN08, Prop. 1.11]).

Definition 1.19. Let $P \subset M_{\mathbb{R}}$ be a n -dimensional Gorenstein polytope of index r and C_P be the corresponding $(n+1)$ -dimensional reflexive Gorenstein cone of index r . We define the *dual n -dimensional Gorenstein polytope* P^* to be the support of the dual reflexive Gorenstein cone $C_P^\vee$, i.e.,

$$C_{P^*} := C_P^\vee.$$

It follows immediately from the definition that $(P^*)^* = P$.

Remark 1.20. It is easy to show that the duality for Gorenstein polytopes commutes with the pyramid construction, i.e., if P and P^* are dual to each other n -dimensional Gorenstein polytopes of index r , then $\Pi(P)$ and $\Pi(P^*)$ are dual to each other $(n+1)$ -dimensional Gorenstein polytopes of index $r+1$. In particular, a Gorenstein polytope P of degree d is a pyramid if and only if the dual Gorenstein polytope P^* is a pyramid.

Theorem 1.21. [BN08, Th. 2.6] *Let $\Delta_1, \dots, \Delta_r \subset M_{\mathbb{R}}$ be lattice polytopes. Assume that the Minkowski sum $\Delta = \Delta_1 + \dots + \Delta_r$ is n -dimensional. Denote by $\tilde{\Delta}$ the Cayley polytope $\Delta_1 * \dots * \Delta_r$ associated with $\Delta_1, \dots, \Delta_r$. Then the following statements are equivalent:*

- (1) $\tilde{\Delta}$ is a $(n+r-1)$ -dimensional Gorenstein polytope of degree n (and index r),
- (2) The polytope Δ is reflexive,
- (3) The cone $C_{\tilde{\Delta}}$ associated with $\tilde{\Delta}$ is a reflexive Gorenstein cone of index r .

This theorem shows a way for constructing examples of Gorenstein polytopes using Minkowski sum decompositions of reflexive polytopes.

Example 1.22. Let Δ be the reflexive cube $[-1, 1]^n \subset \mathbb{R}^n$. Then Δ is a Minkowski sum of $2n$ 1-dimensional lattice polytopes $\Delta_i^\pm := [0, \pm e_i]$ ($i = 1, \dots, n$), where $e_1, \dots, e_n$ the standard basis of $\mathbb{R}^n$. The corresponding $(3n-1)$ -dimensional Gorenstein polytope

$$\Delta_1^+ * \dots * \Delta_n^+ * \Delta_1^- * \dots * \Delta_n^-$$

of degree n will be denoted by Θ_n .

Remark 1.23. In [ET06], I. Emiris and E. Tsigaridas computed all possible Minkowski decompositions of all reflexive lattice polytopes in dimension 2. Their results are listed up to isomorphism in Table 1 and Table 2.

Remark 1.24. Let $P \subset M_{\mathbb{R}}$ be a n -dimensional Gorenstein polytope of index r . By 1.18, the dual polytope P^* of a Gorenstein polytope P of index r is also a Gorenstein polytope of index r . Moreover, P and P^* are combinatorially dual to each other, i.e., there exists a natural bijection $F \leftrightarrow F^*$ between k -dimensional faces F of P and $(n-k-1)$ -dimensional faces F^* of P^* .

	+	+ +
	+	+ +
	+	+ +
	+	+ + +
	+	
	+	
	+	+ +

TABLE 1. Minkowski sum decompositions of reflexive polygons

Remark 1.25. If $r > 1$ then the reflexive polytopes rP and rP^* are not dual to each other as reflexive polytopes (see 1.9).

Denote by $M' \subset M_{\mathbb{R}}$ the n -dimensional lattice generated by M and $m/r \in M_{\mathbb{R}}$, where $m \in rP$ is the unique interior point of the reflexive polytope rP . In order to get the dual Gorenstein polytope P^* one has to dualize the reflexive polytope rP , i.e., to get the dual polytope $(rP)^*$ and replace the dual lattice N by the sublattice $N' \subset N$, $|N/N'| = r$, which is dual to M' (see [BN08, Prop. 1.15]).

Remark 1.26. By [BN08, Prop. 1.16], one has

$$\partial(rP)^* \cap N = \partial(rP)^* \cap N',$$

i.e., there exists a natural bijection between the set of N' -lattice points in P^* and the set of boundary N -lattice points in the dual reflexive polytope $(rP)^*$. Moreover, every facet Γ of P^* considered as a lattice polytope with respect to N' is isomorphic to the facet Γ of Δ^* considered as a lattice polytope with respect to N .



	 + 	
	 + 	
	 + 	
	 + 	
	 + 	 +  + 
	 + 	
	 + 	
	 + 	 +  + 
	 + 	
	 + 	

TABLE 2. Minkowski sum decompositions of reflexive polygons (continued)

Proposition 1.27. *Let $P \subset M_{\mathbb{R}}$ be an n -dimensional Gorenstein polytope of degree d . Denote by $\Gamma_1, \dots, \Gamma_k$ all codimension-1 faces of P . Then one has*

$$\sum_{i=1}^k \text{Vol}_{\mathbb{N}}(\Gamma_i) = (n+1-d) \text{Vol}_{\mathbb{N}}(P).$$

Proof. The polytope $\Delta = (n+1-d)P$ is reflexive. Up to a shift by a lattice vector, we can assume that 0 is the unique interior lattice point of Δ . Then one can decompose Δ into a union of n -dimensional pyramids with vertex 0 over $(n-1)$ -dimensional

faces $(n+1-d)\Gamma_1, \dots, (n+1-d)\Gamma_k$ of Δ . Since heights of all these pyramids are equal to one, we obtain

$$\sum_{i=1}^k \text{Vol}_{\mathbb{N}}((n+1-d)\Gamma_i) = \text{Vol}_{\mathbb{N}}(\Delta).$$

It remains to combine it with the equalities $\text{Vol}_{\mathbb{N}}(\Delta) = (n+1-d)^n \text{Vol}_{\mathbb{N}}(P)$ and

$$\text{Vol}_{\mathbb{N}}((n+1-d)\Gamma_i) = (n+1-d)^{n-1} \text{Vol}_{\mathbb{N}}(\Gamma_i), \quad 1 \leq i \leq k.$$

□

Definition 1.28. Let $v_i \in P$ be a vertex of a n -dimensional Gorenstein polytope P of degree d . We denote by $C(v_i)$ the n -dimensional cone generated by vectors $v - v_i$ where v runs over all lattice points of P .

Proposition 1.29. Let $v_i \in P$ be an arbitrary vertex of a n -dimensional Gorenstein polytope P of degree d and let $\Gamma_i \subset P^*$ be the dual facet in the dual polytope P^* . Then then cone $C(v_i)$ is dual to the cone C_{Γ_i} over the lattice polytope Γ_i (see 1.30).

Example 1.30.




FIGURE 2. Duality between $C(v_1)$ and C_{Γ_1}

Proof. Let $\Delta = (n+1-d)P \subset M_{\mathbb{R}}$ be the reflexive polytope corresponding to P . By 1.25, P^* is obtained from the dual reflexive polytope $\Delta^* \subset N_{\mathbb{R}}$ by choosing a sublattice $N' \subset N$ of index $r = n+1-d$. By 1.26, every facet Γ of P^* with respect to the lattice N' is isomorphic to the facet Γ of Δ^* (with respect to the lattice N). On the other hand, the cone $C(v_i)$ is equal to the cone generated by vectors $v - (n+1-d)v_i$ where v runs over all lattice points of $\Delta = (n+1-d)P$. The duality between reflexive polytopes Δ and Δ^* shows that $C(v_i)$ is dual to cone C_{Γ_i} over Γ_i is a facet of Δ . □

2. THE GRADED GORENSTEIN ALGEBRA S_P

Let K be an arbitrary field of characteristic 0 and let $P \subset M_{\mathbb{R}}$ be an arbitrary n -dimensional lattice polytope. We consider the $(n+1)$ -dimensional lattice $\overline{M} = M \oplus \mathbb{Z}$ and the $(n+1)$ -dimensional cone $C_P \subset \overline{M}_{\mathbb{R}}$ over P . There exists a natural grading of lattice points $\overline{m} = (m, k) \in C_P \cap \overline{M}$ ($m \in M, k \in \mathbb{Z}_{\geq 0}$):

$$\deg \overline{m} := k,$$

so that $C_P \cap \overline{M}$ becomes a graded monoid.

Definition 2.1. We denote by $S_P := K[C_P \cap \overline{M}]$ the graded K -algebra of the semigroup $C_P \cap \overline{M}$. For each lattice point $\overline{m} \in C_P \cap \overline{M}$ we denote by $\mathbf{x}^{\overline{m}}$ the corresponding monomial in S_P . It is well-known that the K -algebra S_P is always Cohen-Macaulay (see e.g. [Sta78]).

Definition 2.2. A lattice point $m \in C_P \cap \overline{M}$ is called *irreducible* if it is not representable as a sum of two lattice points $\overline{m} = \overline{m}' + \overline{m}''$ ($\overline{m}', \overline{m}'' \in C_P \cap \overline{M}$).

Obviously every lattice point of degree 1 in $C_P \cap \overline{M}$ is irreducible. Denote by $I_P(k)$ the number of irreducible lattice points of degree k in $C_P \cap \overline{M}$. Since the semigroup $C_P \cap \overline{M}$ is finitely generated we obtain that

$$I(P, t) := \sum_{k \geq 0} I_P(k) t^k$$

is a polynomial, i.e. $I_P(k) = 0$ for $k \gg 0$. The number $I_P(k)$ can be identified with the number of lattice points of degree k in a minimal generating set of the semigroup $C_P \cap \overline{M}$.

Definition 2.3. Let $S = \bigoplus_{i \geq 0} S_i$ be a finitely generated graded K -algebra, $S_0 \cong K$. For any $k, l \in \mathbb{N}$ we denote by $S_k S_l$ the K -subspace in S_{k+l} generated by products xy , where $x \in S_k$, $y \in S_l$. We define the numbers $g_k(S)$ as follows: $g_0(S) = 1$,

$$g_k(S) := \dim_K S_k / \left(\sum_{j=1}^{k-1} S_j S_{k-j} \right), \quad \forall k > 0.$$

It is clear that $g_1(S) = \dim_K S_1$ and the sum

$$G_S(t) := \sum_{k \geq 0} g_k(S) t^k$$

is a polynomial, i.e. $g_k(S) = 0$ for $k \gg 0$. The number $g_k(S)$ ($k > 0$) can be identified with the number of elements of degree k in the minimal generating system of the K -algebra S . Here the minimality means that any proper subset of this system does not generate S .

Proposition 2.4. *Let $S = S_P$ be the graded algebra of a lattice polytope P . Then the polynomial $G_S(t)$ is equal to $I(P, t)$.*

Proof. Since $S_P = K[C_P \cap \overline{M}]$ is generated by monomials $\mathbf{x}^{\overline{m}}$ corresponding to lattice points $\overline{m} \in C_P \cap \overline{M}$, a minimal generating system $G = \{\gamma_1, \dots, \gamma_l\}$ of the K -algebra S_P can be chosen as a finite subset of these monomials. It remains to show that $\mathbf{x}^{\overline{m}} \in G$ if and only if $\overline{m} \in C_P \cap \overline{M}$ is irreducible. Indeed, if $\mathbf{x}^{\overline{m}} \in G$ and $\overline{m} = \overline{m}' + \overline{m}''$ ($\overline{m}', \overline{m}'' \in C_P \cap \overline{M}$) and $k = \deg \overline{m}'$, $l = \deg \overline{m}''$, then $\mathbf{x}^{\overline{m}} = \mathbf{x}^{\overline{m}'} \mathbf{x}^{\overline{m}''} \in S_k S_l$. This contradicts the minimality of G . On the other hand, if $\overline{m} \in C_P \cap \overline{M}$ is an irreducible lattice point of degree k , then $\mathbf{x}^{\overline{m}}$ can not be a polynomial expression of elements of degree $< k$. Hence $\mathbf{x}^{\overline{m}}$ must appear in G . $\square$

Proposition 2.5. *Let P be a n -dimensional lattice polytope. Denote by A_P an Artinian graded K -algebra obtained by quotient of S_P modulo a regular sequence of $n+1$ elements $y_0, y_1, \dots, y_n$ of degree 1. Then*

$$G_{A_P}(t) = I_P(t) - (n+1)t.$$

Proof. Obviously, we have

$$g_1(A_P) = \dim_K(A_P)_1 = |P \cap M| - (n+1) = I_P(1) - (n+1).$$

Denote by ψ the epimorphism $S_P \rightarrow A_P$. If $G \subset S_P$ is minimal generating system of monomials, then $\psi(G)$ generate A_P as K -algebra. This implies $g_k(A_P) \leq I_P(k) \forall k \geq 2$. In fact, we have the equality $g_k(A_P) = I_P(k) \forall k \geq 2$, because the kernel of the epimorphism

$$\psi : (S_P)_k \rightarrow (A_P)_k$$

is contained in $(S_P)_1(S_P)_{k-1}$. $\square$

Let P be an arbitrary n -dimensional lattice polytope and let $I(P, 1) = \sum_{k \geq 0} I_P(k)$ be the number of irreducible lattice points $C_P \cap \overline{M}$. Then we can describe the graded algebra S_P as a quotient of a polynomial ring $R := K[x_1, \dots, x_{I(P,1)}]$ such that exactly $I_P(k)$ of variables x_i have weight k and $J := \text{Ker}(R \rightarrow S_P)$ is an ideal generated by binomials

$$B = x_{i_1}^{a_1} \cdots x_{i_s}^{a_s} - x_{j_1}^{b_1} \cdots x_{j_r}^{b_r}.$$

We use the following two statements:

Proposition 2.6. [Bat06, Prop. 3.5] *Let $\{B_1, \dots, B_p\}$ be a set of binomials which minimally generate the ideal J . Then P is a pyramid over a $(n-1)$ -dimensional lattice polytope P' , i.e., $P = \Pi(P')$, if and only if there exists a variable $x_i \in \{x_1, \dots, x_{I(P,1)}\}$ which does not appear in any of binomials $B_1, \dots, B_p$.*

Proposition 2.7. [Bat06, Prop. 3.6] *Let $\overline{R} := R/\langle y \rangle$ be a polynomial ring in $I(P, 1) - (n+1)$ variables obtained from the polynomial ring $R := K[x_1, \dots, x_{I(P,1)}]$ by quotient modulo a regular sequence $y_0, y_1, \dots, y_n$ of elements of degree 1. We denote by $\overline{J}$ the ideal $J/\langle y \rangle J$ in $\overline{R}$. Then the numbers and the degrees of minimal generators of ideals $J \subset R$ and $\overline{J} \subset \overline{R}$ are the same.*

Now we apply the above statements to a Gorenstein polytope P . In this case, the graded K -algebras S_P and A_P are known to be Gorenstein. If $A := A_0 \oplus \cdots \oplus A_l$ is an Artinian graded Gorenstein K -algebra ($A_0 = K$, $A_l \neq 0$), then the zero ideal in A is irreducible and the multiplication in A defines a perfect pairing

$$A_i \times A_{l-i} \rightarrow A_l \cong K.$$

For our purposes it will be sufficient to use the following more special fact:

Proposition 2.8. [Hun07] *Let $A := A_0 \oplus A_1 \oplus A_2$ be an Artinian graded K -algebra where $A_0 \cong A_2 \cong K$ is a field, and let $\langle \cdot, \cdot \rangle$ be a symmetric bilinear form on A_1 given by the multiplication*

$$\langle \cdot, \cdot \rangle : A_1 \times A_1 \rightarrow A_2 \cong K \quad (x, y) \mapsto \langle x, y \rangle := xy.$$

Then A is a Gorenstein ring if and only if $\langle \cdot, \cdot \rangle$ is a non-degenerate symmetric bilinear form on A_1 .

This characterisation allows to describe all relations in A if $\dim_K(A_1) \leq 2$:

Proposition 2.9. *Let $A = A_0 \oplus A_1 \oplus A_2$ be an Artinian graded Gorenstein K -algebra and $a = \dim_K(A_1) \in \{0, 1, 2\}$. Then we have:*

- (i) *if $a = 0$, then $A \cong K[X]/(X^2)$, $\deg(X) = 2$;*
- (ii) *if $a = 1$, then $A \cong K[X]/(X^3)$, $\deg(X) = 1$;*
- (ii) *if $a = 2$, then $A \cong K[X, Y]/(XY, X^2 - \lambda Y^2)$, $\deg(X) = \deg(Y) = 1$, $\lambda \neq 0$.*

Proof. (i) Let $x \in A_2$ be a nonzero element. Consider the K -algebra homomorphism $\psi : K[X] \rightarrow A$ defined by the condition $\psi(X) = x$. Since $A_2 A_2 \subset A_4 = 0$ we have $x^2 = 0$, i.e., $X^2 \in \text{Ker } \psi$. So ψ induces an isomorphism $A \cong K[X]/(X^2)$, because both rings A and $K[X]/(X^2)$ are 2-dimensional K -vector spaces.

(ii) Let $x \in A_1$ be a nonzero element. Since the bilinear map $A_1 \times A_1 \rightarrow A_2$ is nondegenerate, we have $x^2 \neq 0$, i.e. ψ is surjective. However, $x^3 = 0$, because $A_3 = 0$. Consider the K -algebra homomorphism $\psi : K[X] \rightarrow A$ defined by the condition $\psi(X) = x$. Then $X^3 \in \text{Ker } \psi$. So ψ induces an isomorphism $R \cong K[X]/(X^3)$, because both rings R and $K[X]/(X^3)$ are 3-dimensional K -vector spaces.

(iii) Since the bilinear map $A_1 \times A_1 \rightarrow A_2 \cong K$ is nondegenerate there exist K -linearly independent elements $x, y \in A_1$ such that $xy = 0$ and $x^2 \neq 0$, $y^2 \neq 0$. Consider the K -algebra homomorphism $\psi : K[X, Y] \rightarrow A$ defined by the conditions $\psi(X) = x$, $\psi(Y) = y$. Then $XY, X^2 - \lambda Y^2 \in \text{Ker } \psi$ for some $\lambda \in K \setminus \{0\}$. Comparing the dimensions of $K[X, Y]/(XY, X^2 - \lambda Y^2)$ and A we get an isomorphism $A \cong K[X, Y]/(XY, X^2 - \lambda Y^2)$. $\square$

Theorem 2.10. *Let P be a n -Gorenstein polytope of degree 2 such that its h^* -vector is $(1, a, 1)$ where $a \in \{0, 1, 2\}$, i.e., $\text{Vol}_{\mathbb{N}}(P) = a + 2 \leq 4$. Assume that P is not a pyramid over a $(n - 1)$ -dimensional Gorenstein polytope. We write the graded Gorenstein ring S_P as quotient of a polynomial ring $K[x_1, \dots, x_l]$ modulo an ideal J generated by binomials. Then $n \leq 5$ and there exist up to isomorphism exactly the following 15 possibilities for P*

- (1) $\dim P = 2$: Δ_i , $1 \leq i \leq 4$;
- (2) $\dim P = 3$: P_i , $1 \leq i \leq 7$;
- (3) $\dim P = 4$: Q_i , $1 \leq i \leq 3$;
- (4) $\dim P = 5$: R_1

The corresponding binomial relations in S_P are presented in the table below

Proof. Let us consider all three values of a separately.

Case 1: $a = 0$. By 2.5, the graded K -algebra S_P is minimally generated by $n + 1$ elements of degree 1 and by one element of degree 2. First we note that $n \geq 3$, because for any 2-dimensional lattice polytope Δ the graded K -algebra S_{Δ} is generated by elements of degree 1. So S_P has $n + 2 \geq 5$ generators. Since $\dim(S_P)_1 = 1$ the polytope P contains exactly $n + 1$ lattice points. These lattice points must be vertices, because

Δ_1	$x_1x_2x_3 = x_4^3$	P_4	$x_1x_2 = x_3^2, x_1x_4 = x_5x_6^2$
Δ_2	$x_1x_2 = x_5^2, x_3x_4 = x_5^2$	P_5	$x_1x_2 = x_3^2, x_3x_4 = x_5x_6$
Δ_3	$x_1x_2 = x_5^2, x_3x_4 = x_1x_5$	P_6	$x_1x_2 = x_3x_4, x_1x_5 = x_2x_6$
Δ_4	$x_1x_2 = x_5^2, x_3x_4 = x_1^2$	P_7	$x_1x_2 = x_5x_6, x_1x_2 = x_3x_4$
P_1	$x_1x_2x_3x_4 = x_5^2$	Q_1	$x_1x_2x_3 = x_4x_5x_6$
P_2	$x_1x_2x_3 = x_4x_5^2$	Q_2	$x_1x_2 = x_3x_4, x_5x_6 = x_7x_1$
P_3	$x_1x_2 = x_3^2, x_4x_5 = x_6^2$	Q_3	$x_1x_2 = x_3x_4, x_5x_6 = x_7^2$
		R_1	$x_1x_2 = x_3x_4, x_5x_6 = x_7x_8$

$\dim P = n$. Thus, P is a n -dimensional lattice simplex. Let $x_1, \dots, x_{n+1}$ be variables in the polynomial ring $R = K[x_1, \dots, x_{n+1}, x_{n+2}]$ whose images in S_P are monomials $\mathbf{x}^{\overline{m}}$ corresponding to vertices $v_1, \dots, v_{n+1}$ of P . Since $\deg x_{n+2} = 2$, the image of x_{n+2} in S_P is a monomial corresponding to an irreducible lattice point $v_{n+1} \in 2P \cap M$ which is not an *integral* nonnegative linear combination of $v_1, \dots, v_{n+1}$. Since P is a convex hull of $v_1, \dots, v_{n+1}$, there exist nonnegative *rational* numbers $c_1, \dots, c_{n+1}$ such that

$$\sum_{i=1}^{n+1} c_i(v_i, 1) = (v_{n+2}, 2).$$

By 2.7, there is only one binomial relation among $n+2$ generators of S_P and this relation has degree 4. This implies, $2c_i \in \mathbb{Z}$ ($i = 1, \dots, n+1$) and the single binomial relation in S_P has form $\prod_{i=1}^{n+1} x_i^{2c_i} = x_{n+2}^2$. Since P is not a pyramid, by 2.6, $2c_i \geq 1$ ($i = 1, \dots, n+1$). On the other hand, $\sum_{i=1}^{n+1} 2c_i = 4 = \deg x_{n+2}^2$. This implies $n+1 \leq 4$. Thus, 3 is the only possible value of n and $2c_1 = 2c_2 = 2c_3 = 2c_4 = 1$. So we come to the binomial relation $x_1x_2x_3x_4 - x_5^2 = 0$. This relation defines a 3-dimensional Gorenstein simplex which we denote by P_1 .


Case 2: $a = 1$. By 2.5 and 2.7, the graded K -algebra S_P is minimally generated by $n+2$ elements of degree 1 satisfying a single cubic binomial relation. Such a binomial relation includes at most 6 variables from $\{x_1, \dots, x_{n+2}\} \subset R$. By 2.6, this implies $n+2 \leq 6$, i.e., $n \leq 4$. Moreover, if $n = 4$, then up to renumeration of variables this binomial must be $x_1x_2x_3 - x_4x_5x_6$ (polytope Q_1). If $n = 2$, then P is a reflexive triangle and the binomial has form $x_1x_2x_3 - x_4^3$ (polytope Δ_1). By 2.6, if $n = 3$ then at least one monomial in the binomial is product of 3 different variables. Without loss of generality, we can assume that this monomial is $x_1x_2x_3$. Then the second monomial must have form $x_i^2x_j$. Up to renumeration of variables we obtain the binomial relation $x_1x_2x_3 - x_4x_5^2$ which defines a 3-dimensional Gorenstein polytope P_2 .

Case 3: $a = 2$. By 2.5 and 2.7, the graded K -algebra S_P is minimally generated by $n+3$ elements of degree 1 satisfying two quadratic binomial relations. These two quadratic relations include at most 8 variables from $\{x_1, \dots, x_{n+3}\} \subset R$. By 2.6, this


implies $n + 3 \leq 8$, i.e., $n \leq 5$. The quadratic monomials appearing in the binomials are either x_i^2 , or $x_i x_j$. It is not difficult to enumerate all possibilities and get the corresponding Gorenstein polytopes

$$\Delta_2, \Delta_3, \Delta_4, P_3, P_4, P_5, P_6, P_7, Q_2, Q_3, R_1.$$


□

 P_1


$$x_1 x_2 x_3 x_4 = x_5^2$$

 P_2


$$x_1 x_2 x_3 = x_4 x_5^2$$

 P_3


$$\begin{aligned} x_1 x_2 &= x_3^2, \\ x_4 x_5 &= x_6^2 \end{aligned}$$

 P_4


$$\begin{aligned} x_1 x_2 &= x_3^2, \\ x_1 x_4 &= x_5 x_6 \end{aligned}$$

 P_5

$$\begin{aligned} x_1 x_2 &= x_3^2, \\ x_3 x_4 &= x_5 x_6 \end{aligned}$$

 P_6

$$\begin{aligned} x_1 x_2 &= x_3 x_4, \\ x_1 x_5 &= x_2 x_6 \end{aligned}$$

 P_7

$$\begin{aligned} x_1 x_2 &= x_5 x_6, \\ x_1 x_2 &= x_3 x_4 \end{aligned}$$

3. GORENSTEIN POLYTOPES OF DEGREE 2

Let $P \subset M_{\mathbb{R}}$ be a n -dimensional Gorenstein polytope of degree 2 with h^* -polynomial $h_P^*(t) = 1 + at + t^2$, $a \geq 0$. Then we have

$$\begin{aligned} i(P, k) &= \binom{n+k}{n} + \binom{n+k-1}{n}a + \binom{n+k-2}{n} \\ &= \frac{1}{k!}(a+2)k^n + \text{terms of lower order}, \\ |P \cap M| = i(P, 1) &= \binom{n+1}{n} + \binom{n}{n}a + \binom{n-1}{n} = n + a + 1. \end{aligned}$$

In particular,

$$\text{Vol}_{\mathbb{N}}(P) = a + 2 = |P \cap M| - n + 1.$$

Proposition 3.1. *Let $P \subset M_{\mathbb{R}}$ be an n -dimensional Gorenstein polytope of degree 2. Then for any lattice subpolytope $P' \subset P$ one has $\text{Vol}_{\mathbb{N}}(P') \leq |P' \cap M| - n + 1$. Moreover, one has:*

- (1) $\text{Vol}_{\mathbb{N}}(P') = |P' \cap M| - n + 1$ if and only if P' is a Gorenstein polytope of degree 2;
- (2) $\text{Vol}_{\mathbb{N}}(P') = |P' \cap M| - n$ if and only if $\deg P' \leq 1$.

Proof. We use the formula for the h^* -polynomial of P

$$h_P^*(t) = 1 + (|P \cap M| - n - 1)t + t^2$$

and the equality $\text{Vol}_{\mathbb{N}}(P) = h_P^*(1) = |P \cap M| - n + 1$. By 1.2, we obtain that either $\deg P' = 2$ and P' is a Gorenstein polytope (see 1.15), or $\deg P' \leq 1$. In the first case, we obtain $\text{Vol}_{\mathbb{N}}(P') = |P' \cap M| - n + 1$ (as for P above). In the second case, the h^* -polynomial of P' is linear and we obtain $\text{Vol}_{\mathbb{N}}(P') = h_{P'}^*(1) = |P' \cap M| - n$. $\square$

Let us make some general remarks about pyramids:

Proposition 3.2. *A n -dimensional Gorenstein polytope P of degree 2 is a pyramid $\Pi(P')$ over an $(n-1)$ -dimensional Gorenstein polytope P' of degree 2 if and only if there exists a proper face $Q \subset P$ such that $\deg Q = 2$.*

Proof. One direction of this statement (i.e., if P is a pyramid) is obvious.

Now we assume that a proper face $Q \subset P$ such that $\deg Q = 2$. Then Q is contained in a facet $\Gamma \subset P$. By 1.2, $\deg Q \leq \deg \Gamma \leq \deg P$ and we conclude that $\deg \Gamma = 2$. Thus, without loss of generality we can assume that Q has codimension 1 (i.e. $Q = \Gamma$). Moreover, since h^* -polynomial is monotone, the leading coefficient of h^* -polynomial of Γ must be 1. Thus, Γ is a Gorenstein polytope of degree 2 (see 1.15). Let p be the unique interior lattice point of $(n-2)\Gamma$. First we show that there exists only one lattice point q of P which is not contained in Γ . If there were two different lattice points $q_1, q_2 \in P \setminus \Gamma$, then we would have two different lattice points $q_1 + p$ and $q_2 + p$ in the interior of $(n-1)P$ (contradiction). Therefore, q is a vertex of P which is combinatorially equivalent to a pyramid over Γ . It remains to show that the

integral height h of the vertex q in this pyramid is 1. One has $\text{Vol}_{\mathbb{N}}(P) = h \text{Vol}_{\mathbb{N}}(\Gamma)$. On the other hand, by 3.1, we have

$$\text{Vol}_{\mathbb{N}}(P) = |P \cap M| - n + 1, \quad \text{Vol}_{\mathbb{N}}(\Gamma) = |\Gamma \cap M| - n.$$

We have already shown that $|P \cap M| = |\Gamma \cap M| + 1$. Therefore, $\text{Vol}_{\mathbb{N}}(P) = \text{Vol}_{\mathbb{N}}(\Gamma)$ and $h = 1$. $\square$

Corollary 3.3. *A Gorenstein polytope P of degree 2 is not a pyramid if and only if all proper faces of P and P^* have degree ≤ 1 .*

Proof. It follows immediately from 1.20 and 3.2. $\square$

From now until the end of this section we assume that $P \subset M_{\mathbb{R}}$ is a 3-dimensional Gorenstein polytope of degree 2.

Proposition 3.4. *If P is a 3-dimensional Gorenstein polytope of degree 2, then*

$$\sum_{E \subset P, \dim E=1} \text{Vol}_{\mathbb{N}}(E) \cdot \text{Vol}_{\mathbb{N}}(E^*) = 12.$$

Proof. Let Δ be a reflexive polytope of dimension 3. Then

$$\sum_{F \subset \Delta, \dim F=1} \text{Vol}_{\mathbb{N}}(F) \cdot \text{Vol}_{\mathbb{N}}(F^*) = 24,$$

where $F^* \subset \Delta$ denote the dual to F 1-dimensional face of the dual reflexive polytope Δ^* . This equality follows from the formula for the Euler number of Calabi-Yau hypersurfaces in toric varieties [BD96, Cor. 7.10] and the fact that the Euler number of $K3$ surfaces is 24 (see also [Haa05]).

We apply this formula to the Gorenstein polytope $\Delta = 2P$. Then $F = 2E$ and hence $\text{Vol}_{\mathbb{N}}(F) = 2 \text{Vol}_{\mathbb{N}}(E)$. On the other hand, by 1.26, the dual 1-dimensional faces F^* and E^* are isomorphic as lattice polytopes, i.e., $\text{Vol}_{\mathbb{N}}(F^*) = \text{Vol}_{\mathbb{N}}(E^*)$. So we get

$$24 = \sum_{F \subset \Delta, \dim F=1} \text{Vol}_{\mathbb{N}}(F) \cdot \text{Vol}_{\mathbb{N}}(F^*) = \sum_{E \subset P, \dim E=1} 2 \text{Vol}_{\mathbb{N}}(E) \cdot \text{Vol}_{\mathbb{N}}(E^*).$$

Dividing by 2, we obtain the required equality. $\square$

Example 3.5. Let Δ be a 2-dimensional reflexive polytope and Δ^* its dual. By 1.20, $P := \Pi(\Delta)$ and $P^* = \Pi(\Delta^*)$ are two dual to each other 3-dimensional Gorenstein polytopes of degree 2. A 1-dimensional face $E \subset P$ is either an edge connecting the vertex of the pyramid $\Pi(\Delta)$ to a vertex $v_i \in \Delta$ ($\text{Vol}_{\mathbb{N}}(E) = 1$), or a 1-dimensional face of $\Gamma_j \subset \Delta$. Then the dual 1-dimensional face $E^* \subset P^*$ is respectively either 1-dimensional face $v_i^* \subset \Delta^*$, or an edge connecting the vertex of the pyramid $\Pi(\Delta^*)$ with dual vertex $\Gamma_j^* \in \Delta^*$ ($\text{Vol}_{\mathbb{N}}(E^*) = 1$). Therefore

$$\sum_{E \subset P, \dim E=1} \text{Vol}_{\mathbb{N}}(E) \cdot \text{Vol}_{\mathbb{N}}(E^*) = \sum_i 1 \cdot \text{Vol}_{\mathbb{N}}(v_i^*) + \sum_j \text{Vol}_{\mathbb{N}}(\Gamma_j) \cdot 1.$$

By 1.27, we have

$$\text{Vol}_{\mathbb{N}}(\Delta) = \sum_j \text{Vol}_{\mathbb{N}}(\Gamma_j), \quad \text{Vol}_{\mathbb{N}}(\Delta^*) = \sum_i \text{Vol}_{\mathbb{N}}(v_i^*).$$

So for 3-dimensional Gorenstein polytopes $P := \Pi(\Delta)$ and $P^* = \Pi(\Delta^*)$ the equality in 3.4 is equivalent to the well-known identity for reflexive polygons:

$$\text{Vol}_{\mathbb{N}}(\Delta) + \text{Vol}_{\mathbb{N}}(\Delta^*) = 12.$$

Example 3.6. Another example is shown in Figure 3 below. The Gorenstein simplex P has 6 edges $E_1, \dots, E_6$ ($\text{Vol}_{\mathbb{N}}(E_i) = 1$). Its dual polytope P^* has also 6 edges $E_1^*, \dots, E_6^*$ ($\text{Vol}_{\mathbb{N}}(E_i^*) = 2$). Therefore, we have

$$\sum_{E \subset P, \dim E=1} \text{Vol}_{\mathbb{N}}(E) \cdot \text{Vol}_{\mathbb{N}}(E^*) = \sum_{i=1}^6 \text{Vol}_{\mathbb{N}}(E_i) \cdot \text{Vol}_{\mathbb{N}}(E_i^*) = 6 \cdot (1 \cdot 2) = 12.$$




FIGURE 3. Dual Gorenstein polytopes in dimension 3

For any face $F \subset P$ we denote by $\text{Int}(E)$ the number of lattice points in the relative interior of F .

Proposition 3.7. *Let P be a 3-dimensional Gorenstein polytope of degree 2 which is not a pyramid. Then*

$$|P \cap M| + |P^* \cap N| + \sum_{E \subset P} \text{Int}(E) \cdot \text{Int}(E^*) = 14.$$

Proof. If P is not a pyramid, then all lattice points of P are contained in edges (see 3.3), i.e.,

$$|P \cap M| = v + \sum_{E \subset P} \text{Int}(E),$$

where v denotes the number of vertices of P . Denote by e and f the numbers of edges and faces of P respectively. By Euler formula, $e = v + f - 2$. Applying 3.4 and

$\text{Vol}_{\mathbb{N}}(E) = \text{Int}(E) + 1$, one has

$$\begin{aligned}
12 &= \sum_{i=1}^e (\text{Int}(E_i) + 1)(\text{Int}(E_i^*) + 1) \\
&= e + \sum_{i=1}^e (\text{Int}(E_i) + \text{Int}(E_i^*) + \text{Int}(E_i) \cdot \text{Int}(E_i^*)) \\
&= v + f - 2 + \sum_{i=1}^e \text{Int}(E_i) + \sum_{i=1}^e \text{Int}(E_i^*) + \sum_{i=1}^e \text{Int}(E_i) \cdot \text{Int}(E_i^*).
\end{aligned}$$

Using $|P \cap M| = \sum_{i=1}^e \text{Int}(E_i) + v$ and $|P^* \cap N| = \sum_{i=1}^e \text{Int}(E_i^*) + f$, we obtain the statement. $\square$

Example 3.8. The dual to each other polytopes P and P^* in Example 3.6 contain respectively 4 and 10 lattice points. Since all lattice points of P are vertices, so we obtain $\sum_{E \subset P} \text{Int}(E) \cdot \text{Int}(E^*) = 0$, i.e.,

$$|P \cap M| + |P^* \cap N| + \sum_{E \subset P} \text{Int}(E) \cdot \text{Int}(E^*) = 4 + 10 + 0 = 14.$$

Proposition 3.9. *If P is a 3-dimensional Gorenstein polytope of degree 2 which is not a pyramid, then*

$$\text{Vol}_{\mathbb{N}}(P) + \text{Vol}_{\mathbb{N}}(P^*) \leq 10.$$

Proof. Recall that $\text{Vol}_{\mathbb{N}}(P) = |P \cap M| - 2$ and $\text{Vol}_{\mathbb{N}}(P^*) = |P^* \cap N| - 2$. By 3.7 and by the obvious inequality

$$\sum_{E \subset P} \text{Int}(E) \cdot \text{Int}(E^*) \geq 0,$$

we obtain the statement. $\square$

4. MINIMAL GORENSTEIN POLYTOPES AND THE CLASSIFICATION

Definition 4.1. Let $P \subset M_{\mathbb{R}}$ be an arbitrary n -dimensional Gorenstein polytope of degree 2 with vertices $\{v_1, \dots, v_k\}$. We call P *minimal* if for all $j \in \{1, \dots, k\}$ the convex hull of the set $\{v_1, \dots, v_k\} \setminus \{v_j\}$ is not a Gorenstein polytope of degree 2. Analogously, we call P *maximal*, if for any lattice point $v_{k+1} \in M \setminus \{v_1, \dots, v_k\}$ the convex hull of the lattice points $\{v_1, \dots, v_k, v_{k+1}\}$ is not Gorenstein polytope of degree 2.

Remark 4.2. If a n -dimensional Gorenstein polytope P of degree 2 is isomorphic to a subpolytope of another n -dimensional Gorenstein polytope Q of degree 2, then we write $P \preceq Q$. If $P \subseteq Q$, then we immediately obtain that both reflexive polytopes $(n-1)P$ and $(n-1)Q$ contain a *common* unique interior lattice point $m \in M$. By 1.25, we obtain that the dual Gorenstein polytope Q^* is naturally contained in the

dual polytope P^* , i.e. $Q^* \preceq P^*$. Therefore the duality $P \leftrightarrow P^*$ establishes a bijection between the set of minimal and the set of maximal Gorenstein polytopes:

$$\{\text{minimal Gorenstein polytopes}\} \leftrightarrow \{\text{maximal Gorenstein polytopes}\}.$$

Proposition 4.3. *Let $v_i \in P$ be an arbitrary vertex of a n -dimensional Gorenstein polytope P of degree 2 and let $\Gamma_i^* \subset P^*$ be the facet of the dual polytope P^* which is dual to v_i . Then*

$$P_i := \text{conv}(\{P \cap M\} \setminus \{v_i\})$$

is a Gorenstein polytope of degree 2 if and only if the facet Γ_i^ is a standard $(n-1)$ -dimensional basic simplex. In particular, P is minimal if and only if $\text{Vol}_{\mathbb{N}}(\Gamma^*) \geq 2$ for all facets $\Gamma^* \subset P^*$.*

Proof. By 3.1, we obtain that P_i is Gorenstein of degree 2 if and only if

$$\text{Vol}_{\mathbb{N}}(P_i) = |P_i \cap M| - n + 1 = |P \cap M| - n - 2 = \text{Vol}_{\mathbb{N}}(P) - 1.$$

The latter holds true if and only if P_i is obtained from P by cutting out a n -dimensional lattice polytope S_i with $\text{Vol}_{\mathbb{N}}(S_i) = 1$ (i.e., S_i is a basic simplex). In this case, the simplex S_i is a convex hull of $v_i \in P$ and a simplicial facet $\Gamma_i \subset P_i$, where $\text{Vol}_{\mathbb{N}}(\Gamma_i) = 1$, i.e., Γ_i is a standard $(n-1)$ -dimensional basic simplex. $\square$

Corollary 4.4. *If P and Q two n -dimensional Gorenstein polytopes of degree 2 such that $Q \preceq P$. Then*

$$\text{Vol}_{\mathbb{N}}(P) + \text{Vol}_{\mathbb{N}}(P^*) = \text{Vol}_{\mathbb{N}}(Q) + \text{Vol}_{\mathbb{N}}(Q^*).$$

Proof. We use the induction on $k = |P \cap M| - |Q \cap M|$. If $k = 0$ then $P = Q$ and there is nothing to prove. If $k \geq 1$ then $P \neq Q$ and there exists a vertex $v_i \in P$ which is not contained in Q . Let $P_i := \text{conv}(\{P \cap M\} \setminus \{v_i\})$. Then $Q \preceq P_i$. Moreover, the proof of 4.3 shows that P_i is obtained from P by cutting out a n -dimensional lattice basic simplex $S_i = \text{conv}(\Gamma_i, v_i)$. Let v_i^* be the dual vertex of P_i corresponding to the facet $\Gamma_i \subset P_i$. Then

$$P^* := \text{conv}(\{P_i^* \cap N\} \setminus \{v_i^*\}).$$

As in the proof of 4.3, we obtain

$$\text{Vol}_{\mathbb{N}}(P^*) = \text{Vol}_{\mathbb{N}}(P_i^*) - 1.$$

Together with $\text{Vol}_{\mathbb{N}}(P_i) = \text{Vol}_{\mathbb{N}}(P) - 1$ this implies

$$\text{Vol}_{\mathbb{N}}(P) + \text{Vol}_{\mathbb{N}}(P^*) = \text{Vol}_{\mathbb{N}}(P_i) + \text{Vol}_{\mathbb{N}}(P_i^*).$$

Since $Q \preceq P_i$ and $|P_i \cap M| - |Q \cap M| < |P \cap M| - |Q \cap M|$ we can apply the induction hypothesis and get

$$\text{Vol}_{\mathbb{N}}(P_i) + \text{Vol}_{\mathbb{N}}(P_i^*) = \text{Vol}_{\mathbb{N}}(Q) + \text{Vol}_{\mathbb{N}}(Q^*).$$

This proves the statement. $\square$

The main purpose of this section is to prove the following result:

Theorem 4.5. *Let P be a minimal n -dimensional Gorenstein polytope of degree 2. Then $\text{Vol}_{\mathbb{N}}(P) \leq 4$.*

Remark 4.6. If P is a pyramid $\Pi(P')$ over a $(n-1)$ -dimensional Gorenstein polytope P' of degree 2, then $\text{Vol}_{\mathbb{N}}(P) = \text{Vol}_{\mathbb{N}}(P')$ and minimality of P is equivalent to minimality of P' . Therefore it is sufficient to prove 4.5 for Gorenstein polytopes of degree 2 which are not pyramids.

Lemma 4.7. *Let P be a n -dimensional Gorenstein polytope of degree 2 and P^* its dual. Assume that*

$$\text{Vol}_{\mathbb{N}}(P^*) \leq 4.$$

Then either P is not minimal, or $\text{Vol}_{\mathbb{N}}(P) \leq 4$.

Proof. We can apply the classification of all n -dimensional Gorenstein polytopes P^* of degree 2 satisfying the condition $\text{Vol}_{\mathbb{N}}(P^*) \leq 4$ (see 2.10).

If $\text{Vol}_{\mathbb{N}}(P^*) \in \{2, 3\}$ a direct check via this classification shows that all facets of P^* are standard simplices. Thus, by 4.3, we obtain that $P = (P^*)^*$ is not minimal.

If $\text{Vol}_{\mathbb{N}}(P^*) = 4$, then P^* may contain facets which are not standard simplices, but if $\text{Vol}_{\mathbb{N}}(P) \geq 5$ there always exists at least one facet which is a standard simplex. By 4.3, this implies that P is not minimal. $\square$

By Lemma 4.7, the statement of Theorem 4.5 follows from:

Theorem 4.8. *Let P be a minimal n -dimensional Gorenstein polytope of degree 2. Then at least one of two inequalities $\text{Vol}_{\mathbb{N}}(P) \leq 4$ and $\text{Vol}_{\mathbb{N}}(P^*) \leq 4$ holds.*

We prove Theorem 4.8 in several steps:

Let P be a minimal n -dimensional Gorenstein polytope of degree 2. We fix a vertex $v_i \in P$ and the facet $\Gamma_i^* \subset P^*$ which is dual to v_i . Without loss of generality we assume that $v_i = 0$, i.e., $C(v_i) \supset P$ is a cone with vertex 0.

Since P^* is not a pyramid, by Corollary 3.3, we obtain that Γ_i^* is a $(n-1)$ -dimensional Gorenstein polytope of degree ≤ 1 . By 4.3, the facet Γ_i^* can not have degree 0, i.e., Γ_i^* is not a $(n-1)$ -dimensional basic lattice simplex. Therefore, we have $\deg \Gamma_i^* = 1$. By the complete classification of lattice polytopes of degree 1 (see 1.5), Γ_i^* is either an $(n-1)$ -dimensional exceptional simplex, or a Lawrence prism with heights $h_1, \dots, h_{n-1}$ such that $\text{Vol}_{\mathbb{N}}(\Gamma_i^*) = \sum_{j=1}^{n-1} h_j \geq 2$.

Lemma 4.9. *If Γ_i^* is a $(n-1)$ -dimensional exceptional simplex, then $\text{Vol}_{\mathbb{N}}(P) = 2$.*

Proof. Using 1.29, one obtains that the cone $C(v_i) \supset P$ is isomorphic to the simplicial cone generated by $2e_1 - e_2 - e_3, e_2, e_3, \dots, e_n$, where $e_1, \dots, e_n$ is a basis of M . Then $n+1$ lattice points $0, 2e_1 - e_2 - e_3, e_2, \dots, e_n$ are contained in P and their convex hull P'_i is already a simplex whose h^* -polynomial is $1 + t^2$. By minimality, we have $P = P'_i$. Therefore $\text{Vol}_{\mathbb{N}}(P) = \text{Vol}_{\mathbb{N}}(P'_i) = 2$. $\square$

Lemma 4.10. *If Γ_i^* is a $(n-1)$ -dimensional Lawrence prism with heights $h_1, \dots, h_{n-1}$ such that $\text{Vol}_{\mathbb{N}}(\Gamma_i^*) = \sum_i h_i \geq 3$, then $\text{Vol}_{\mathbb{N}}(P) \in \{3, 4\}$.*

Proof. Using 1.29, one obtains that the cone $C(v_i) \supset P$ is isomorphic to the cone generated by $n + 1$ vectors

$$e_1, \dots, e_n, -e_n + \sum_{i=1}^{n-1} h_i e_i,$$

where $e_1, \dots, e_n$ is a basis of M . We denote by P'_i the convex hull of 0 and $n + 1$ lattice points $e_1, \dots, e_n, -e_n + \sum_{i=1}^{n-1} h_i e_i$. Then a direct calculation shows that $\text{Vol}_{\mathbb{N}}(P'_i) = \sum_{i=1}^{n-1} h_i$. Since P'_i lies between two parallel hyperplanes $z_n = \pm 1$ and the intersections of P'_i with these hyperplanes are exactly two vertices $p := e_n$ and $q := -e_n + \sum_{i=1}^{n-1} h_i e_i$, all remaining lattice points of P'_i are contained in the hyperplane $\{z_n = 0\}$. We want to show that the set $M \cap P' \cap \{z_n = 0\}$ is either $\{0, e_1, \dots, e_{n-1}\}$, or it possibly has one more lattice point $(p + q)/2$ (if $(p + q)/2 \in M$). Indeed, by 3.3 all lattice points of P are contained in 1-dimensional faces of P . Since $P'_i \subset P$, the same is true also for P'_i , i.e. all lattice points of P'_i are contained in 1-dimensional faces of P'_i . But P'_i is a convex hull of $n + 2$ lattice points and all segments connecting two of them (except maybe $[p, q]$) do not have interior lattice points, because the convex hull of $\{0, e_1, \dots, e_{n-1}\}$ is a basic $(n - 1)$ -dimensional simplex in the hyperplane $\{z_n = 0\}$ and the z_n -coordinates of p and q are 1 and -1 respectively. We note that $\{z_n = 0\} \cap [p, q] = (p + q)/2 = (\sum_{i=1}^{n-1} h_i e_i)/2$. Thus, we obtain $|P'_i \cap M| \leq n + 3$. By 3.1, the latter implies $\text{Vol}_{\mathbb{N}}(P'_i) \leq |P'_i \cap M| - n + 1 \leq 4$, i.e. $\sum_{i=1}^{n-1} h_i \in \{3, 4\}$. If $\sum_{i=1}^{n-1} h_i = 3$, then $(p + q)/2 \notin M$ can not be a lattice point of P'_i and $|P'_i \cap M| = n + 2$. Together with $\text{Vol}_{\mathbb{N}}(P'_i) = 3$ this implies that P'_i is a Gorenstein polytope with h^* -polynomial $1 + t + t^2$. If $\sum_{i=1}^{n-1} h_i = 4$, then by 3.1, P'_i must be a Gorenstein polytope of degree 2, i.e., $|P'_i \cap M| = n + 3$, $(p + q)/2$ is a lattice point of P'_i , and $\text{Vol}_{\mathbb{N}}(P'_i) = 4$. By minimality, we have in both cases $P = P'_i$, i.e. $\text{Vol}_{\mathbb{N}}(P) \in \{3, 4\}$. $\square$

The following statement completes the proof of Theorem 4.8 and Theorem 4.5:

Lemma 4.11. *Assume that P is a minimal Gorenstein polytope of degree 2 such that every codimension-1 face $\Gamma_i^* \subset P^*$ is a Lawrence prism with $\text{Vol}_{\mathbb{N}}(\Gamma_i^*) = 2$. Then either $\text{Vol}_{\mathbb{N}}(P) \leq 4$, or $\text{Vol}_{\mathbb{N}}(P^*) \leq 4$.*

Proof. In this situation, all generators of the cone $C(v_i)$ are contained in an affine hyperplane H_i which divides P into two lattice polytopes $P_i = \text{conv}(\{P \cap M\} \setminus \{v_i\})$ and a Lawrence prism P'_i containing the vertex v_i such that $\text{Vol}_{\mathbb{N}}(P'_i) = 2$. Moreover, P'_i is a pyramid with vertex v_i over the $(n - 1)$ -dimensional Lawrence prism $\overline{P}_i = H_i \cap P$.

By minimality, P_i must have degree 1 and by 1.5 we have the following two cases:

CASE 1: The polytope P_i is an exceptional simplex, i.e., $\text{Vol}_{\mathbb{N}}(P_i) = 4$. Then its facet $\overline{P}_i$ is also a simplex. Moreover, P'_i must be also a simplex, since P'_i is a pyramid over $\overline{P}_i$. Let k be the number of vertices of P (this is also the number of facets in P^*). The whole polytope P is a union of simplices P_i and P'_i . Therefore $k \leq n + 2$.

By 1.27, we have

$$(n-1) \operatorname{Vol}_{\mathbb{N}}(P^*) = \sum_{i=1}^k \operatorname{Vol}_{\mathbb{N}}(\Gamma_i^*) \leq 2(n+2).$$

If $n \geq 4$, then we obtain $\operatorname{Vol}_{\mathbb{N}}(P^*) \leq 2 + 6/(n-1) \leq 4$. If $n = 3$, then $\operatorname{Vol}_{\mathbb{N}}(P) = \operatorname{Vol}_{\mathbb{N}}(P'_i) + \operatorname{Vol}_{\mathbb{N}}(P_i) = 2 + 4 = 6$. By 3.9, we have $\operatorname{Vol}_{\mathbb{N}}(P^*) \leq 4$.

CASE 2: The polytope P_i is a Lawrence prism. Since P'_i and $\overline{P}_i$ are both Lawrence prisms and $\operatorname{Vol}_{\mathbb{N}}(P'_i) = \operatorname{Vol}_{\mathbb{N}}(\overline{P}_i) = 2$ the number of vertices of $\overline{P}_i$ is at most $n+1$. On the other hand, $\overline{P}_i$ is a facet of P_i which is not a basic $(n-1)$ -dimensional simplex. Therefore P_i contains at most 2 more vertices than in $\overline{P}_i$. Together with $v_i \in P'_i$ we see that $P = P'_i \cup P_i$ contains $k \leq n+4$ vertices. By 1.27, we have

$$(n-1) \operatorname{Vol}_{\mathbb{N}}(P^*) = \sum_{i=1}^k \operatorname{Vol}_{\mathbb{N}}(\Gamma_i^*) \leq 2(n+4).$$

If $n \geq 5$, then we obtain $\operatorname{Vol}_{\mathbb{N}}(P^*) \leq 2 + 10/(n-1) < 5$, i.e. $\operatorname{Vol}_{\mathbb{N}}(P^*) \leq 4$.

If $n = 4$, then $(n-1) \operatorname{Vol}_{\mathbb{N}}(P^*) = \sum_{i=1}^k \operatorname{Vol}_{\mathbb{N}}(\Gamma_i^*)$ together with $\operatorname{Vol}_{\mathbb{N}}(\Gamma_i^*) = 2$ implies that the number k of vertices of P must be divisible by 3. Since $k \leq n+4 = 8$, we obtain $k = 6$ and $\operatorname{Vol}_{\mathbb{N}}(P^*) = 4$.

If $n = 3$, then it remains to exclude only the case $\operatorname{Vol}_{\mathbb{N}}(P) = \operatorname{Vol}_{\mathbb{N}}(P^*) = 5$ (all other cases follow from 3.9). By 1.27, in the last case both polytopes P and P^* must have 5 vertices. There exist exactly two different combinatorial types of 3-dimensional polytopes P with 5 vertices, but only for one of these types the dual polytope P^* has also 5 vertices. Therefore, both polytopes P and P^* must be combinatorially equivalent to a pyramid over a lattice 4-gon S such that $\operatorname{Vol}_{\mathbb{N}}(S) = 2$. This implies that $\operatorname{Vol}_{\mathbb{N}}(P) = \operatorname{Vol}_{\mathbb{N}}(P^*) = 5$ is divisible by $\operatorname{Vol}_{\mathbb{N}}(S) = 2$. Contradiction. $\square$

Proposition 4.12. *Every n -dimensional Gorenstein polytope $P \neq 2S_3$ of degree 2 is a Cayley polytope $P = \Delta_1 * \dots * \Delta_r$ where $\Delta_1, \dots, \Delta_r$ are plane lattice polytopes such that the Minkowski sum $\Delta = \Delta_1 + \dots + \Delta_r$ is a reflexive polygon.*

Proof. It is enough to assume that P is not a pyramid, because a pyramid over a Cayley polytope is again a Cayley polytope. On the other hand, every n -dimensional subpolytope of a Cayley polytope $\Delta_1 * \dots * \Delta_r$ is again a Cayley polytope $\Delta'_1 * \dots * \Delta'_r$, where Δ'_i is a lattice subpolytope of Δ_i ($i = 1, \dots, r$). By 4.5, if P is a maximal Gorenstein polytope of degree 2, then $\operatorname{Vol}_{\mathbb{N}}(P^*) \leq 4$, i.e., P^* is one of polytopes classified in 2.10. One can check the whole list of these Gorenstein polytopes P^* and obtain that among potential maximal polytopes P only the 3-dimensional Gorenstein polytope $2S_3$ can not be described as a Cayley polytope. Moreover, if v_i is a vertex of $2S_3$ and P_i is the convex hull of $(2S_3 \cap M) \setminus \{v_i\}$ then P_i is again a Cayley polytope. This shows that any proper Gorenstein subpolytope of $2S_3$ is also a Cayley polytope. $\square$

Theorem 4.13. *There exist exactly 37 n -dimensional Gorenstein polytopes of degree 2 which are not pyramids:*

- (1) 16 *reflexive polygons*;
- (2) 3-dimensional Gorenstein polytopes $P_1, \dots, P_{15}$;
- (3) 4-dimensional Gorenstein polytopes $Q_1, \dots, Q_5$;
- (4) 5-dimensional Gorenstein polytope R_1 .


Proof. There are two ways to prove this statement.

The first one is to use 4.12 and the classification of all possible Minkowski sum decompositions of reflexive polygons due to Emiris and Tsigaridas [ET06].


The second one is to enumerate all Gorenstein subpolytopes of degree 2 of maximal Gorenstein polytopes of degree 2. The maximal Gorenstein polytopes are contained in the list of all P such that $a_{P^*} = \text{Vol}_{\mathbb{N}}(P^*) - 2 \leq 2$ (see 2.10).

Both ways do not need any computer and can be easily realized. $\square$


Below we give a complete list of n -dimensional ($n \geq 3$) Gorenstein polytopes P of degree 2 which are not pyramids. This list includes some of combinatorial invariants of P such as its f -vector $f = (f_0, \dots, f_n)$ (where f_i denotes the number of i -dimensional faces of P), the dual partner P^* , the lattice normalized volume $\text{Vol}_{\mathbb{N}}(P)$ as well as all possible descriptions of P as a Cayley polytope.




$$\begin{array}{c}
 P_1 \quad \text{---} \text{---} * \text{---} \text{---} \\
 \hline
 f = (1, 4, 6, 4, 1) \\
 P_1^* = P_{15} \\
 \text{Vol}_{\mathbb{N}}(P_1) = 2
 \end{array}$$




$$\begin{array}{c}
 P_2 \quad \text{---} \text{---} * \text{---} \text{---} \\
 \hline
 f = (1, 5, 9, 6, 1) \\
 P_2^* = P_{14} \\
 \text{Vol}_{\mathbb{N}}(P_2) = 3
 \end{array}$$



$$\begin{array}{c}
 P_3 \quad \text{---} * \text{---} \text{---} \text{---} \\
 \hline
 f = (1, 4, 6, 4, 1) \\
 P_3^* = P_3 \\
 \text{Vol}_{\mathbb{N}}(P_3) = 4
 \end{array}$$




$$\begin{array}{c}
 P_4 \quad \text{---} * \text{---} \text{---} \text{---} \\
 \hline
 f = (1, 5, 8, 5, 1) \\
 P_4^* = P_{13} \\
 \text{Vol}_{\mathbb{N}}(P_4) = 4
 \end{array}$$




$$P_5 \quad \begin{array}{c} \triangle \\ \vdots \\ * \quad \downarrow \end{array}$$

$$\begin{array}{l} f = (1, 5, 9, 6, 1) \\ P_5^* = P_{10} \\ \text{Vol}_{\mathbb{N}}(P_5) = 4 \end{array}$$




$$P_6 \quad \begin{array}{c} \square \quad * \\ \vdots \quad \downarrow \\ \triangle \quad * \quad \triangle \end{array}$$

$$\begin{array}{l} f = (1, 6, 11, 7, 1) \\ P_6^* = P_{11} \\ \text{Vol}_{\mathbb{N}}(P_6) = 4 \end{array}$$




$$P_7 \quad \begin{array}{c} \triangle \quad * \quad \triangle \\ \vdots \end{array}$$

$$\begin{array}{l} f = (1, 6, 12, 8, 1) \\ P_7^* = P_{12} \\ \text{Vol}_{\mathbb{N}}(P_7) = 4 \end{array}$$




$$P_8 \quad \begin{array}{c} \square \quad * \quad \triangle \\ \vdots \end{array}$$

$$\begin{array}{l} f = (1, 7, 12, 7, 1) \\ P_8^* = P_8 \\ \text{Vol}_{\mathbb{N}}(P_8) = 5 \end{array}$$











$$P_9 \quad \begin{array}{c} \triangle \quad * \quad \triangle \\ \vdots \end{array}$$

$$\begin{array}{l} f = (1, 6, 10, 6, 1) \\ P_9^* = P_9 \\ \text{Vol}_{\mathbb{N}}(P_9) = 5 \end{array}$$



$$P_{10} \quad \begin{array}{c} \triangle \quad * \quad \triangle \\ \vdots \quad \downarrow \\ \square \quad * \quad \downarrow \end{array}$$

$$\begin{array}{l} f = (1, 6, 9, 5, 1) \\ P_{10}^* = P_5 \\ \text{Vol}_{\mathbb{N}}(P_{10}) = 6 \end{array}$$

	
P_{11}  $f = (1, 7, 11, 6, 1)$ $P_{11}^* = P_6$ $\text{Vol}_{\mathbb{N}}(P_{11}) = 6$	P_{12}  $f = (1, 8, 12, 6, 1)$ $P_{12}^* = P_7$ $\text{Vol}_{\mathbb{N}}(P_{12}) = 6$
	
P_{13}  $f = (1, 5, 8, 5, 1)$ $P_{13}^* = P_4$ $\text{Vol}_{\mathbb{N}}(P_{13}) = 6$	P_{14}  $f = (1, 6, 9, 5, 1)$ $P_{14}^* = P_2$ $\text{Vol}_{\mathbb{N}}(P_{14}) = 7$
	P_{15} none $f = (1, 4, 6, 4, 1)$ $P_{15}^* = P_1$ $\text{Vol}_{\mathbb{N}}(P_{15}) = 8$

The list of all 4-dimensional Gorenstein non-pyramids Q_i of index 2:

$ \begin{array}{c} Q_1 \quad \begin{array}{c} \bullet \diagdown \bullet \\ \bullet \text{---} \bullet \end{array} * \begin{array}{c} \bullet \text{---} \bullet \end{array} * \begin{array}{c} \updownarrow \end{array} \\ \hline f = (1, 6, 15, 18, 9, 1) \\ Q_1^* = Q_4 \\ \text{Vol}_{\mathbb{N}}(Q_1) = 3 \end{array} $	$ \begin{array}{c} Q_2 \quad \begin{array}{c} \bullet \text{---} \bullet \end{array} * \begin{array}{c} \updownarrow \end{array} * \begin{array}{c} \updownarrow \end{array} \\ \hline f = (1, 7, 17, 18, 8, 1) \\ Q_2^* = Q_5 \\ \text{Vol}_{\mathbb{N}}(Q_2) = 4 \end{array} $
$ \begin{array}{c} Q_3 \quad \begin{array}{c} \bullet \text{---} \bullet \end{array} * \begin{array}{c} \updownarrow \end{array} * \begin{array}{c} \updownarrow \end{array} \\ \hline f = (1, 6, 13, 13, 6, 1) \\ Q_3^* = Q_3 \\ \text{Vol}_{\mathbb{N}}(Q_3) = 4 \end{array} $	$ \begin{array}{c} Q_4 \quad \begin{array}{c} \bullet \diagdown \bullet \\ \bullet \text{---} \bullet \end{array} * \begin{array}{c} \bullet \diagdown \bullet \\ \bullet \text{---} \bullet \end{array} * \begin{array}{c} \bullet \diagdown \bullet \\ \bullet \text{---} \bullet \end{array} \\ \hline f = (1, 9, 18, 15, 6, 1) \\ Q_4^* = Q_1 \\ \text{Vol}_{\mathbb{N}}(Q_4) = 6 \end{array} $

$$\begin{array}{c}
 Q_5 \quad \begin{array}{c} \begin{array}{c} \updownarrow \end{array} * \begin{array}{c} \bullet \diagdown \bullet \\ \bullet \text{---} \bullet \end{array} * \begin{array}{c} \bullet \diagdown \bullet \\ \bullet \text{---} \bullet \end{array} \\
 \begin{array}{c} \updownarrow \end{array} * \begin{array}{c} \bullet \text{---} \bullet \end{array} * \begin{array}{c} \bullet \diagdown \bullet \\ \bullet \text{---} \bullet \end{array} \end{array} \\
 \hline
 f = (1, 8, 18, 17, 7, 1) \\
 Q_5^* = Q_2 \\
 \text{Vol}_{\mathbb{N}}(Q_5) = 5
 \end{array}$$

There exists a single 5-dimensional Gorenstein (selfdual) non-pyramid R_1 of index 3:

$$\begin{array}{c}
 R_1 \quad \begin{array}{c} \bullet \text{---} \bullet \end{array} * \begin{array}{c} \bullet \text{---} \bullet \end{array} * \begin{array}{c} \updownarrow \end{array} * \begin{array}{c} \updownarrow \end{array} \\
 \hline
 f = (1, 8, 24, 34, 24, 8, 1) \\
 R_1^* = R_1 \\
 \text{Vol}_{\mathbb{N}}(R_1) = 4
 \end{array}$$

5. TORIC ALMOST DEL PEZZO MANIFOLDS

A smooth n -dimensional toric variety X is called *almost Del Pezzo manifold* if the anticanonical class $-K_X$ is a semiample Cartier divisor which defines a K_X -trivial birational morphism $\varphi : X \rightarrow X'$, where X' is a Gorenstein toric Del Pezzo variety. Our classification of Gorenstein polytopes P of degree 2 in 4.13 is equivalent to a biregular classification of all n -dimensional Gorenstein toric Del Pezzo varieties, i.e., of all X' . Let P be a n -dimensional Gorenstein polytope of degree 2. We denote by X_P the corresponding toric variety. First we need to understand when a given n -dimensional Gorenstein toric Del Pezzo variety X_P admits a smooth crepant resolution.

Theorem 5.1. *Let P a n -dimensional Gorenstein polytope of degree 2 which is not a pyramid over a $(n - 1)$ -dimensional Gorenstein polytope. Then the Gorenstein toric Del Pezzo variety X_P always admits a crepant desingularization $\widehat{X}_P$.*

Proof. Since a projective toric variety is smooth if and only if all its 0-dimensional torus orbits are nonsingular points, it is enough to analyse the singularities of 0-dimensional torus orbits in X_P which correspond to vertices of the Gorenstein polytope P . If $v_i \in P$ such a vertex, then, by 1.29, the cone $C(v_i)$ generated by $v - v_i$ ($v \in P$) is dual to the cone C_{Γ_i} over the dual facet $\Gamma_i \subset P^*$. By 3.3, all facets of P^* are lattice polytopes of degree ≤ 1 . In [BN07, Prop. 5.1] it was shown that every lattice polytope of degree ≤ 1 admits a unimodular triangulation. This implies existence of smooth crepant resolutions of all 0-dimensional orbits in X_P . $\square$

Example 5.2. The statement in 5.1 is not true if P is a pyramid. Let $X_{\Pi(P_{15})}$ be 4-dimensional Gorenstein toric Del Pezzo variety which is a cone over the 3-dimensional Gorenstein toric Del Pezzo variety $X_{P_{15}}$. Then the vertex of the cone is a terminal quotient singularity $\mathbb{A}^4 / \pm id$, which does not admit a crepant resolution.

Remark 5.3. A smooth crepant desingularization of a Gorenstein toric Del Pezzo variety is not uniquely determined, because there might be many different unimodular triangulations of lattice polytopes of degree 1. For example, 3-dimensional Gorenstein toric Del Pezzo variety $X_{P_{13}}$ admits exactly two different crepant desingularizations related by a flop, because all except one facets of the dual polytope $P_4 = P_{13}^*$ admit a unique unimodular triangulation and the exceptional face defines an isolated conifold singularity in $X_{P_{13}}$, which has exactly two different crepant resolutions. Another example of a 3-dimensional Gorenstein toric Del Pezzo variety with many different crepant resolutions is a 3-dimensional singular cubic $X_{P_2} = \{x_1x_2x_3 - x_4x_5^2 = 0\} \subset \mathbb{P}_4$.

Theorem 5.4. *Let P be a n -dimensional Gorenstein polytope of degree 2 and let $(1, a_{P^*}, 1)$ be the h^* -vector of the dual Gorenstein polytope P^* . Then the Picard number of the crepant resolution $\widehat{X}_P$ is equal to $a_{P^*} + 1$. In particular, if X_P corresponds to a maximal Gorenstein polytope P , then $\text{rk Pic}(\widehat{X}_P) \leq 3$.*

Proof. Let V be a smooth projective n -dimensional toric variety defined by a fan Σ . Denote by $\Sigma^{(1)}$ the set of all 1-dimensional cones in Σ . The Picard number of V equals $|\Sigma^{(1)}| - n$ [Oda88]. Consider now the case $V = \widehat{X}_P$. Then the fan Σ defining V is a simplicial subdivision of the fan defining the Gorenstein toric Del Pezzo variety X_P . By [Bat94], the latter is the fan of cones over proper faces of the dual reflexive polytope $((n-1)P)^*$, i.e., $|\Sigma^{(1)}| = \partial((n-1)P)^* \cap N$. By 1.26, the number of N -lattice points in $\partial((n-1)P)^*$ equals the number of N' -lattice points in the dual Gorenstein polytope P^* . The latter is equal to $a_{P^*} + n + 1$ (see Section 3). So we obtain $\text{rk Pic}(\widehat{X}_P) = a_{P^*} + 1$. The last statement follows from the duality of Gorenstein polytopes and from the inequality $\text{Vol}_{\mathbb{N}}(P^*) = 2 + a_{P^*} \leq 4$ (see 4.5). $\square$

Theorem 5.5. *Let P and Q be two n -dimensional Gorenstein polytopes of degree 2 which are not pyramids such that $Q \preceq P$. Then, up to finitely many flops, the almost Del Pezzo manifold $\widehat{X}_Q$ is obtained from the almost Del Pezzo manifold $\widehat{X}_P$ by blow ups of $\text{Vol}_{\mathbb{N}}(P) - \text{Vol}_{\mathbb{N}}(Q)$ smooth points.*

Proof. Consider Q as a lattice subpolytope of P . Let $v_i \in P$ be a vertex of P which is not contained in Q . In the proof of 4.3 it was shown that $P_i := \text{conv}(\{M \cap P\} \setminus \{v_i\})$ is another Gorenstein polytope of degree 2 containing Q . Moreover, P_i is obtained from P by cutting out a n -dimensional basic simplex S_i ($\text{Vol}_{\mathbb{N}}(S_i) = 1$) containing v_i as one of its vertices, i.e., we have $\text{Vol}_{\mathbb{N}}(P) - \text{Vol}_{\mathbb{N}}(P_i) = 1$ and the vertex v corresponds to a smooth fixed point $p(v_i) \in X_P$. Let Γ_i be the facet of S which does not contain v_i . Then Γ_i is also a facet of P_i and it defines a torus invariant divisor in X_{P_i} which is isomorphic to $\mathbb{P}^{n-1}$, because $\Gamma_i \subset S_i$ is a $(n-1)$ -dimensional basic simplex. By 4.4, we have

$$1 = \text{Vol}_{\mathbb{N}}(P) - \text{Vol}_{\mathbb{N}}(P_i) = \text{Vol}_{\mathbb{N}}(P_i^*) - \text{Vol}_{\mathbb{N}}(P^*) = a_{P_i^*} - a_{P^*}.$$

Using 5.4, we obtain

$$\text{rk Pic } \widehat{X}_{P_i} - \text{rk Pic } \widehat{X}_P = 1.$$

Choose an unimodular triangulation of facets of P^* defining a crepant birational morphism $\widehat{X}_P \rightarrow X_P$. We can extend this triangulation to an unimodular triangulation of faces of $P_i^* \supset P^*$ which defines a crepant birational morphism $\widehat{X}_{P_i} \rightarrow X_{P_i}$. The compatibility of triangulations defines a birational morphism $\varphi : \widehat{X}_{P_i} \rightarrow \widehat{X}_P$ which contracts a divisor $D \cong \mathbb{P}^{n-1} \subset \widehat{X}_{P_i}$ to a smooth point $p(v_i)$. Thus, $\widehat{X}_{P_i}$ is a blow up of a smooth point in $\widehat{X}_P$. We can repeat the same procedure $\text{Vol}_{\mathbb{N}}(P) - \text{Vol}_{\mathbb{N}}(Q)$ times and obtain the statement. $\square$

Theorems 5.1, 5.4, 5.5 show that our combinatorial method in the biregular classification of all Gorenstein toric Del Pezzo varieties is parallel to the birational method in the classification of almost Del Pezzo manifolds due to Jahnke and Peternell [JP08].

REFERENCES

- [Bat82] V. V. Batyrev, *Boundedness of the degree of multidimensional toric Fano varieties*. Vestnik Moskov. Univ. Ser. I Mat. Mekh. (1982), no. 1, 22–27.
- [Bat85] V. V. Batyrev, *Higher dimensional toric varieties with ample anticanonical class*, PhD thesis, 1985. (in russian)
- [Bat94] V. V. Batyrev, *Dual polyhedra and mirror symmetry for Calabi-Yau hypersurfaces in toric varieties*. J. Algebraic Geom. **3** (1994), no. 3, 493–535.
- [BD96] V. V. Batyrev, D.I. Dais, *Strong McKay correspondence, string-theoretic Hodge numbers and mirror symmetry*. Topology 35 (1996), no. 4, 901–929.
- [BB97] V.V. Batyrev, L. A. Borisov, *Dual cones and mirror symmetry for generalized Calabi-Yau manifolds*. in Mirror symmetry, II, (1997), 71–86.
- [Bat06] V. V. Batyrev, *Lattice polytopes with a given h^* -polynomial*. in Algebraic and geometric combinatorics, Contemp. Math., **423** (2006), 1–10.
- [BN07] V.V. Batyrev, B. Nill, *Multiples of lattice polytopes without interior lattice points*. Mosc. Math. J. **7** (2007), 195–207.

- [BN08] V.V. Batyrev, B. Nill, *Combinatorial Aspects of Mirror Symmetry*, Contemp. Math., **452** (2008), 35–66.
- [BR07] W. Bruns, T. Römer, *h-vectors of Gorenstein polytopes*. J. Combin. Theory Ser. A **114** (2007), no. 1, 65–76.
- [CK99] D. A. Cox, S. Katz, *Mirror symmetry and algebraic geometry*. Mathematical Surveys and Monographs, **68**. American Mathematical Society, Providence, RI, 1999.
- [Dem80] M. Demazure, *Surfaces de del Pezzo* In: Séminaire Sur Les Singularités Des Surfaces, Lecture Notes in Mathematics, **777** (1980), 23–69.
- [DuV34] P. Du Val, *On isolated singularities of surfaces which do not affect the conditions of adjunction. I* Proc. Camb. Philos. Soc. **30** (1934), 453–459.
- [Ehr67] E. Ehrhart, *Sur un problème géométrie diophantienne linéaire II* J. Reine Angew. Math. **227** (1967), 25–49.
- [ET06] I.Z. Emiris, E.P. Tsigaridas, *Minkowski decomposition of convex lattice polygons*. in Algebraic geometry and geometric modeling, Math. Vis., Springer, Berlin, (2006) 217–236.
- [Fuj80] T. Fujita, *On the structure of polarized manifolds with total deficiency one I*, J. Math. Soc. Japan **32** (1980), 709–725.
- [Fuj81] T. Fujita, *On the structure of polarized manifolds with total deficiency one II* J. Math. Soc. Japan **33** (1981), 415–434.
- [Fuj84] T. Fujita, *On the structure of polarized manifolds with total deficiency one III*, J. Math. Soc. Japan **36** (1984), 75–89.
- [Fuj90a] T. Fujita, *Classification theories of polarized varieties*, London Mathematical Society Lecture Note Series, **155**. Cambridge University Press, Cambridge, 1990.
- [Fuj90b] T. Fujita, *On singular Del Pezzo varieties*. Algebraic geometry (L’Aquila, 1988), Lecture Notes in Math., **1417** (1990), 117–128,
- [Ful93] W. Fulton, *Introduction to Toric Varieties*, Annals of Mathematics Studies **131**, 1993.
- [Haa05] C. Haase, *Problems from the Cottonwood Room*. in “Integer points in polyhedra—geometry, number theory, algebra, optimization”, Contemp. Math. **374** (2005), 179–191.
- [HNP08] C. Haase, B. Nill, S. Payne, *Cayley decompositions of lattice polytopes and upper bounds for h^* -polynomials*. Preprint arXiv:0804.3667.
- [Hib92] T. Hibi, *Dual polytopes of rational convex polytopes*, Combinatorica **12** (1992), no. 2, 237–240.
- [HS02] L. Hille, H. Skarke, *Reflexive polytopes in dimension 2 and certain relations in $SL_2(\mathbb{Z})$* , J. Algebra Appl. **1** (2002), no. 2, 159–173.
- [Hun07] C. Huneke, *Lectures on local cohomology*, (with an Appendix by Amelia Taylor), in “Interactions between homotopy theory and algebra”, Contemp. Math. **436** (2007), 51–99.
- [Isk77] V. A. Iskovskikh. *Fano 3-folds I*. Math. USSR Izv. **11** (1977), 485–527.
- [Isk78] V. A. Iskovskikh. *Fano 3-folds II*. Math. USSR Izv. **12** (1978), 469–506.
- [Isk80] V. A. Iskovskikh. *Anticanonical models of three-dimensional algebraic varieties*, Current problems in mathematics, Vol. **12** (1980), 59–157. (in russian)
- [JP08] P. Jahnke, T. Peternell, *Almost Del Pezzo Manifolds*, Adv. Geom. **8** (2008), no. 3, 387–411.
- [Koe91] R. J. Koelman, *The Number of Moduli of Families of Curves on Toric Surfaces*, PhD thesis, 1991.
- [KS98] M. Kreuzer, H. Skarke, *Classification of reflexive polyhedra in three dimensions*. Adv. Theor. Math. Phys. **2** (1998), no. 4, 853–871.

- [KS00] M. Kreuzer, H. Skarke, *Complete classification of reflexive polyhedra in four dimensions*. Adv. Theor. Math. Phys. **4** (2000), no. 6, 1209–1230.
- [Nil08] B. Nill, *Lattice polytopes having h^* -polynomials with given degree and linear coefficient*, European J. Combin. **29** (2008), no. 7, 1596–1602.
- [Oda88] T. Oda, *Convex bodies and algebraic geometry. An introduction to the theory of toric varieties*, Ergebnisse der Mathematik und ihrer Grenzgebiete **15**, Springer-Verlag, Berlin, 1988.
- [PR00] B. Poonen, F. Rodriguez-Villegas, *Lattice polygons and the number 12*. Amer. Math. Monthly **107** (2000), no. 3, 238–250.
- [Rab89] S. Rabinowitz, *A census of convex lattice polygons with at most one interior lattice point*. Ars Combin. **28** (1989), 83–96.
- [RST05] D. Repovsh, M. Skopenkov, M. Tsentsel, *An elementary proof of the twelve lattice point theorem*, Math. Zamet. **77** (2005), no. 1-2, 108–111.
- [Sta78] R. Stanley, *Hilbert functions of graded algebras*, Adv. in Math. **28** (1978), 57–83.
- [Sta80] R. Stanley, *Decompositions of rational convex polytopes*, Annals of Discrete Math. **6** (1980), 333–342.

MATHEMATISCHES INSTITUT, UNIVERSITÄT TÜBINGEN, AUF DER MORGENSTELLE 10, 72076 TÜBINGEN, GERMANY

E-mail address: victor.batyrev@uni-tuebingen.de

MATHEMATISCHES INSTITUT, UNIVERSITÄT TÜBINGEN, AUF DER MORGENSTELLE 10, 72076 TÜBINGEN, GERMANY

E-mail address: dorothee@juny.de